


QVA
P 933
1846

THE
PREScriBER'S
PHARMACOPÆIA



THE
✓ PRESCRIBER'S
PHARMACOPÆIA:

CONTAINING
ALL THE MEDICINES

IN

The London Pharmacopœia,

ARRANGED IN CLASSES ACCORDING TO THEIR ACTION, WITH
THEIR COMPOSITION AND DOSES.


BY A PRACTISING PHYSICIAN.

Altered to correspond with the U. S. Dispensatory.

REVISED AND IMPROVED
BY AN AMERICAN PHYSICIAN.

SECOND AMERICAN FROM THE THIRD LONDON EDITION.

NEW YORK:
SAMUEL S. & WILLIAM WOOD,
No. 261, Pearl-street.



QVA
Pq33
1846

Film 8122, item 6

Entered, according to Act of Congress, in the Year 1846,

BY S. S. AND W. WOOD,

In the Clerk's Office of the District Court, for the Southern
District of New-York.

TO

JOHN CONOLLY, M.D.

RESIDENT PHYSICIAN OF THE HANWELL ASYLUM,

In sincere admiration and grateful acknowledgement of his noble and successful efforts to ameliorate the condition of the poor Lunatic, by the substitution of a rational system of moral discipline for mechanical restraint, and parental care and kindness for violence and brute force, this little Work is, on public grounds, inscribed

BY

THE AUTHOR.

ADVERTISEMENT

TO THE THIRD EDITION.

THE singular favour shown by the profession to this little work, evinced by the rapid sale of two large editions, makes it especially incumbent on the Editor to lose no opportunity of adding to its unpretending merits.

The present Edition has been carefully revised. Some errors found to exist in the former have been corrected: and several not unimportant additions have been made in various parts of it. It is therefore hoped that, in the condition which it now appears, it will be found still more deserving the patronage of those for whom it is intended.

DEC. 15, 1844.

ADVERTISEMENT

TO THE SECOND EDITION.

THE very favourable reception which this little work has received from the profession, evinced by the exhaustion of a large impression within the year, has satisfactorily proved the justness of the author's opinion, that it was wanted, and would be useful. It is hoped that the present Edition, improved as it is in several respects, will be found still more completely to fulfil its object. In preparing it for the press, care has been taken to correct and amend some defects which were discovered on a close examination. Numerous additions have been made, and several changes have been adopted in the plan and arrangement, which will render the work still more useful. The principal of these are the following :—

1. The arrangement of the different CLASSES of Medicines in alphabetical order, as well as the individual articles in each class.
2. The separation, under a distinct head, of the external and chirurgical remedies from those for internal use.
3. The addition, in an Appendix, of many useful Tables.
4. The reduction of all the weights and measures (with one or two trifling exceptions) to the same denomination, or at least to the same class of signs, viz, grains, scruples, drachms, ounces, (gr., ℥, ℥, ℥); the measures of pint and gallon (O. cong.) being now rendered in ounces.

It has been thought unnecessary to prefix the *f* (*fluid*) to the ℥ and ℥ when representing *measured* quantities, as the omission can lead to no mistake, every one knowing that solids are weighed, and fluids measured.

N.B. In stating the *composition* of medicines, the exact *quantities* given in the Pharmacopœia have not always been retained, but the *proportions* only.

Nov 1, 1842.

ADVERTISEMENT

TO THE FIRST EDITION.

As this little work had its origin in a want daily experienced by the compiler in his practice, he believes that it will be useful to a numerous class of practitioners who possess a memory of the same general character as his own. He has always had great difficulty in calling to mind, at will, a number of heterogeneous particulars which had not been originally contemplated in a systematic order, although he might be well acquainted with them individually. This difficulty he has often painfully experienced when attempting to pass in review the various drugs and officinal formulæ suited to fulfil any special indication that presented itself in the treatment of a disease: and he has frequently witnessed, in the practice of others, a barrenness of prescription and a consequent confined range of

resources, originating in the same cause, which could not fail to be injurious to the patient and the reputation of the physician. It is expected that the habit of viewing the articles of the *Materia Medica* in some such order as that adopted in the present compilation, will tend, in some degree at least to remedy these evils.

It is further believed that practitioners, from forgetfulness of the officinal preparations, are often led to prescribe extemporaneous formulæ while there exist better substitutes in the *Pharmacopœia*; and it is hoped that the marshalling together all the various compound forms of medicine under the title of the main ingredient, as in these tables, will at once facilitate and simplify the labours of the prescriber.

It is not pretended that the classification of medicines here adopted is even moderately accurate. One perfectly accurate is impossible. All that is intended or hoped for is an arrangement that will allow such a division of the substances as may be at once simple and practically useful. No attempt at scientific refinement in classification has been made. All the older and more common divisions and terms have been retained, as likely to be best understood and best remembered; and no effort has been made

to prevent the repetition of the same medicines in different sections of the classification.

Perhaps in a work of such slight pretensions as this, it is hardly necessary to advert to the sources whence its materials are derived. It may, however, be as well to state that wherever the Pharmacopeia and his own knowledge failed him, the author confidently turned to the inexhaustible store provided in Dr. Pereira's Elements of Materia Medica. In minor matters he has also benefited by the excellent compendium of Dr. Lane.

It will, perhaps, be remarked that the doses of medicines are, in many instances, below the usual standard—at least in their minor term. This is not unintentional; as the author is of opinion that the general habit of practitioners in this country is to prescribe medicines in doses not merely unnecessarily but injuriously large.

MAY, 1841.

ADVERTISEMENT

TO THE

SECOND AMERICAN EDITION.

THE present edition, from the third London, has been carefully compared with the 6th edition of the U. S. Dispensatory by Wood and Bache, and the formulæ altered to correspond with those given in it. In some instances the formulæ of the London Dispensatory have been retained, where new ones have not been ordered in our standard. The *uses* of medicines have been introduced to supply what appeared a defect in the London edition.

AMERICAN EDITOR.

CONTENTS.

PART I.

Remedies for Internal Use.

	PAGE
ACIDS (ANTALKALINES)	17
ALKALINE REMEDIES (ANTACIDS)	ib.
ALTERATIVES, DEOBSTRUENTS	21
ANTHELMINTICS	26
ANTISPASMODICS	27
AROMATICS, AROMATIC STIMULANTS	31
AROMATIC STIMULANT BITTERS	36
ASTRINGENTS	39
CARMINATIVES	45
DIAPHORETICS	49
DILUENTS, DEMULCENTS	52
DIURETICS	55
EMETICS	61
EMMENAGOGUES	63
ERRHINES	65
EXPECTORANTS	ib.
NARCOTICS, SEDATIVES	67
PURGATIVES	76
REFRIGERANTS	89
SIALAGOGUES, MASTICATORIES	92
STIMULANTS	ib

	PAGE
TONICS (MINERAL)	101
TONICS (VEGETABLE)	105

PART II.

For External and Local Use.

CATAPLASMATA (POULTICES)	110
CAUSTICA (ESCHAROTICA)	111
CERATA (OINTMENTS)	113
EMPLASTRA (PLASTERS)	116
ENEMATA (INJECTIONS)	117
METHODUS ENDERMICUS	118
GARGARISMATA (GARGLES)	ib.
LINIMENTA (LINIMENTS)	119
PULVERES (POWDERS)	122
UNGUENTA (OINTMENTS)	113
INHALATION	122

PART III.

POISONS AND COUNTERPOISONS	123
TEMPERATURE OF BATHS	123
PROPORTION OF OPIUM, &c.	129
DOSES ACCORDING TO AGE	130
MINERAL WATERS	131
SALINE DRAUGHTS	135
INDEX	137

WEIGHTS AND MEASURES

USED IN MEDICINE.

WEIGHTS.

Pound.	Ounce.	Drachm.	Scruple.	Grain.
1	= 12	= 96	= 288	= 5760
	1	= 8	= 24	= 480
		1	= 3	= 60
			1	= 20

MEASURES.

Gall.	Pint.	Ounce.	Drachm.	Minim.	Imp. grains.
1	= 8	= 160	= 1280	= 76800	= 70000
	1	= 20	= 160	= 9600	= 8750
		1	= 8	= 480	= 437.5
			1	= 60	= 54.7
				1	= 0.91

AMMONIA.

1. LIQUOR AMMONIÆ. (Aqua Ammoniæ.) (STIMULANTS.)

Dose. ℥ v–xx, (in watery vehicles.)

2. AMMONIÆ SESQUICARBONAS. (Olim A. SUBCARBONAS.)

Dose. Gr. ij–x, in pill or liquid. (STIMULANTS.)

LIQUOR AMMONIÆ SESQUICARBONATIS.

Comp. Am. sesquicarb. ℥ iv, water f ℥ xx. (STIMULANTS.)

Dose. f ℥ ss–j.

- | | |
|--|---------------|
| a. <i>Spiritus Ammoniæ.</i> | } STIMULANTS. |
| b. <i>Spiritus Ammoniæ Aromaticus.</i> | |
| c. <i>Spiritus Ammoniæ Fætidus.</i> | |
- ANTISPASMODICS.

CALX. (Calx recens. Rarely used in medicine.)

1. LIQUOR CALCIS. (Lime water.)

Comp. A saturated solution. 11½ grs. in the pint of water at 60°.

Dose. f ℥ ij–vj, (with milk.)

2. Creta Præparata. (Chalk purified by water.)

Dose. Gr. x–3 j.

3. PULVIS CRETÆ COMPOSITUS.

Comp. 12 chalk, 8 cinnamon, 6 tormentil, 6 acacia, 1 long pepper.

Dose. Gr. x–xx.

Use. In diarrhoea with acidity and without inflammatory symptoms.

4. PULVIS CRETÆ COMPOSITUS CUM OPIO.

Comp. 39 pulv. cret. comp. (No. 3,) 1 opium.

Dose. Gr. v- $\frac{3}{4}$ ss. ($\frac{1}{4}$ gr. opium in gr. x.)

Use. Same as above, but more efficacious.

5. MISTURA CRETÆ.

Comp. Chalk $\frac{3}{4}$ ss, sugar 3 ij, gum arabic 3 ij,
cinnamon water and water $\ddot{\text{aa}}$ f $\frac{3}{4}$ iv.

Dose. f $\frac{3}{4}$ ss-ij.

6. TESTÆ PRÆPARATÆ. (Prepared oyster shells.)

Dose. Gr. x-3 j.

CONFECTIO AROMATICA. (See AROMATICS.)

MAGNESIA.

1. MAGNESIÆ CARBONAS. (Olim SUBCARBONAS.
Magnesia alba.) (PURGATIVES.)

2. MAGNESIA. (M. usta. Calcined M.)

Dose. Gr. v-xx (in aromatic water or milk.)
Better as antacid than No. 1. (PURGATIVES.)

Use. As an antacid in dyspepsia, sick headache,
gout. For children. For excess of uric acid.

POTASSA.

1. LIQUOR POTASSÆ.

Comp. Carb. potassæ $\frac{1}{2}$ j, lime $\frac{1}{2}$ ss, water 1 gall.

Use. In lepra, psoriasis and other cutaneous disorders. To correct acid state of urine.

Dose. \mathfrak{M} x-xxx. (In infus. aurant., beer, broth.)

[BRANDISH'S ALKALINE SOLUTION, is made by adding to a gallon of boiling water, pearlashes, $\frac{3}{4}$ xij, quicklime and woodashes, of each $\frac{3}{4}$ iv. It was given by the inventor in doses of 3 j to 3 iij in beer twice a-day.]

2. POTASSÆ CARBONAS. (Potassæ subcarb.)

Dose. Gr. v-x-xxx.

3. POTASSÆ BICARBONAS. (Potassæ carb.)

Dose. Gr. v-x-xxx.

4. LIQUOR POTASSÆ CARBONATIS.

Comp. Carb. pot. $\frac{1}{2}$ j, water f $\frac{2}{3}$ xij.*Dose.* Mx-3j.

5. LIQUOR POTASSÆ EFFERVESCENS.

Comp. Bicarb. pot. 3j, water f $\frac{2}{3}$ xx, carb. acid,
q. s.*Dose.* f $\frac{2}{3}$ iv, ad libitum.

SAPO. (Hard Soap. Soda and olive oil.)

Dose. Gr. v-xx (chiefly in combination.)

SODA.

Use. To correct too acid urine.

1. SODÆ CARBONAS. (Olim SODÆ SUBCARB.)

Dose. Gr. v-xxx.2. SODÆ CARBONAS EXSICCATA. (Deprived of
water of crystallization.)*Dose.* Gr. iij-xv.

3. SODÆ SESQUICARBONAS. (Ol. SODÆ CARB.)

Dose. Gr. x-3j.

4. LIQUOR SODÆ EFFERVESCENS. (Soda water.)

Comp. Sesquicarb sod. 3j, water f $\frac{2}{3}$ xx, carbonic
acid q. s.*Dose.* f $\frac{2}{3}$ iv, or ad libitum. (Better made by add-
ing carb. sodæ (3j to f $\frac{2}{3}$ xx) to the common
bottle soda water.)

Alteratives. Deobstruents.

ANTIMONIUM? See EMETICS and DIAPHORETICS.)

ARSENICUM. (See TONICS.)

BARII CHLORIDUM. (Muriate of baryta.)

1. LIQUOR BARII CHLORIDI.

Comp. Bar. chlor. $\frac{3}{2}$ j, water f $\frac{3}{2}$ iiij.

Dose. $\frac{1}{2}$ v.

Use. In scrofula, cancer, painful ulcers, cutaneous eruptions.

BROMINIUM. (Bromine.)

Dose and Form. Gr. 1-16 to gr. ss, dissolved in water. (Rarely used.)

1. POTASSII BROMIDUM.

Dose. Gr. j-x, in pill or solution.

Use. In scrofulous tumours, bronchocele.

CALCII CHLORIDUM. (Muriate of Lime.)

(See TONICS.)

1. LIQUOR CALCII CHLORIDI.

Comp. Marble in fragments $\frac{3}{2}$ ix, acid hydrochl. Oj, aq. distill. q. s.

Dose. f $\frac{3}{2}$ ss-f $\frac{3}{2}$ j.

Use. Tabes mesenterica, scrofulous affections.

GUAIACUM. (See DIAPHORETICS.)

Use. Secondary syphilis, cutaneous eruptions.

HYDRARGYRUM. (Mercury.) (See PURGATIVES.)

1. HYDRARGYRUM CUM CRETâ.

Comp. 3 mercury, 5 chalk

Dose. Gr. ij-v.

Use. Principally for children: in deficient biliary secretion.

2. PILULÆ HYDRARGYRI.

Comp. Mercury $\frac{3}{j}$, confect. roses $\frac{3}{j}$ iss, liquorice $\frac{3}{j}$ ss.

Dose. Gr. j-v.

Use. A mild mercurial.

3. HYDRARGYRI OXYDUM.

Dose. Gr. ss-ij. (Rarely used.)

4. HYDRARGYRI CHLORIDUM. (Submuriæ. Calomel.)

Dose. Gr. $\frac{1}{4}$ -j.

5. PILULÆ HYDRARGYRI CHLORIDI COMPOSITÆ. (Plummer's Pill.)

Comp. 1 calomel, 1 sulphuret of antimony, 1 treacle, 2 guaiacum, (1 gr. calomel in v.)

Dose. Gr. iij-v-x.

Use. Chronic rheumatism. Scaly eruptive diseases.

6. HYDRARGYRI BICHLORIDUM. (Oxymuriæ. Corrosive sublimate.)

Dose. Gr. $\frac{1}{16}$ - $\frac{1}{8}$.

Use. Secondary syphilis, cutaneous diseases.

7. LIQUOR HYDRARGYRI BICHLORIDI.

Comp. Bichlor. of merc., hydrochlorate of ammonia, $\ddot{\text{aa}}$ gr. x; water f $\frac{3}{j}$ xx.

Dose. $\frac{3}{ss}$ -ij in mild liquids, e. g. linseed tea. (Gr. ss. of hyd. bi. in $\frac{3}{j}$, gr. 1-16 in $\frac{3}{j}$.)

8. HYDRARGYRI IODIDUM.

Comp. Mercury $\frac{3}{j}$, iodine $\frac{3}{v}$, alcohol q. s.

Dose. Gr. j-ijj.

Use. In scrofula and scrofulous syphilis.

9. HYDRARGYRI BINIODIDUM. (More energetic than No. 8.)

Comp. Cor. sub. 3*i*, iod. pot. 3*x*, aq. dist. Oij.

Dose. Gr. 1-16 - $\frac{1}{4}$.

Use. Same as above.

10. PILULÆ HYDRARGYRI IODIDI.

Comp. 1 iodide of mercury, 1 ginger, 3 confec-
tion of rose. (1 gr. of iod. of merc. in gr. v.)

Dose. Gr. v-x.

11. HYDRARGYRI BISULPHURETUM. (Cinnabar.
Red sulphuret.)

Dose. (Rarely used internally.) For fumigation,
3ss.

12. HYDRARGYRI SULPHURETUM CUM SULPHURE.
(Black sulph., Æthiops mineral.)

Dose. Gr. v-xx.

13. UNGUENTUM HYDRARGYRI FORTIUS. (See
PART II.)

14. UNGUENTUM HYDRARGYRI MITIUS. (See
PART II.)

15. CERATUM HYDRARGYRI COMPOSITUM. (See
PART II.)

16. LINIMENTUM HYDRARGYRI COMPOSITUM. (See
PART II.)

17. EMPLASTRUM HYDRARGYRI. (See PART II.)

18. EMPLASTRUM AMMONIACI CUM HYDRARGYRO.
(See PART II.)

19. UNGUENTUM HYDRARGYRI NITRATIS. (See
PART II.)

20. UNGUENTUM HYDRARGYRI NITRICO-OXYDI-
 (See PART II.)
21. UNGUENTUM HYDRARGYRI IODIDI. (See
 PART II.)
22. UNGUENTUM HYDRARGYRI BINIODIDI. (See
 PART II.)

IODINIUM. (Iodine.)

Dose. Gr. ss-ij, in pills—for immediate use.
 (Rarely used in substance.)

1. TINCTURA IODINII COMPOSITA.

Comp. Iod. $\frac{3}{2}$ ss, iod. of pot. $\frac{3}{2}$ j, alc. Oj.

Dose. \mathfrak{M} v- \mathfrak{Z} j, (dilute when used.)

Use. Mostly externally. Erysipelas, chilblains.

2. IODIDUM HYDRARGYRI. (See HYDRARG.)

**3. BINIODIDUM HYDRARGYRI. (See HYDRARGY-
 RUM.)**

4. POTASSII IODIDUM. (Hydriodas Potassæ.)

Dose. Gr. j-v-x- \mathfrak{D} j.

Use. Scrofula, syphilitic rheumatism, strumous ophthalmia, secondary syphilis.

5. LIQUOR POTASSII IODIDI COMPOSITUS.

Comp. Iod. of potassium $\frac{3}{2}$ iss, iodine $\frac{3}{2}$ vj, water Oj.

Dose. \mathfrak{M} vj, containing iodine gr. $\frac{1}{4}$.

6. PLUMBI IODIDUM.

Dose. Gr. $\frac{1}{4}$ -ij.

Use. For discussion of scrofulous tumours.

7. UNGUENTUM PLUMBI IODIDI. (See PART II.)

8. UNGUENTUM IODINII COMPOSITUM. (See PART II.)

9. UNGUENTUM HYDRARGYRI IODIDI. (See PART II.)

10. UNGUENTUM HYDRARGYRI BINIODIDI. (See PART II.)

11. FERRI IODIDUM. (Iodide of iron.)

Dose. Gr. ij-x, in distilled water, or thick colourless syrup.

SARZA. (Smilax officinalis. Sarsaparilla. The root.)

Dose. 3ss-ij, in powder.

1. DECOCTUM SARZÆ.

Comp. Sarza 3 ijss, water 3 xl, boil to 3 xx.

Dose. f 3 iv-viij. (Usually with extract or syrup.)

2. DECOCTUM SARZÆ COMPOSITUM. (Lisbon diet drink.)

Comp. Sarsap. 3 vj, sassaf. guaiacum and liquorice each 3 j, mezereon 3 iij, aq. Oiv.

Dose. f 3 iv-f 3 vj, ter in die.

Use. In secondary syphilis.

3. SYRUPUS SARZÆ.

Comp. Decoct. sarzæ and sugar.

Dose. f 3 iij-3 j. (Of feeble power.)

4. EXTRACTUM SARZÆ. The watery extract.

Dose. 3j-3ij. (Rubbed down with water, flavoured with tinct. aurant. or volatile oil:—also made into pills.)

5. SYR. SARSAP. COMP. U. S.

Comp. Sarsap. 1bij, guaiac. 3 iij, rose leaves, senna and liquorice root, each 3 ij, ol. sassaf. and anisi, each 1v, ol. gaulth. 1lij, dil. alcohol Ox, sugar 1b viij.

Dose. f 3 ss.

Use. In syphilis, cutaneous eruptions, scrofula.

Anthelmintics.

ASPIDIUM. (Filix mas. The male Fern. The root.)

Dose. 3 j-ij, in powder.

[**NOTE.**—Nouffleur's plan:—3 ij in water ʒ iv; after six hours, a bolus of calomel (gr. x), scammony and Gamboge.]

Use. In cases of tænia.

GRANATUM. (Punica granatum. Pomegranate. Bark of the Root.)

Dose. Decoction (ʒ j of fresh root in water f ʒ xx, boil to f ʒ x) f ʒ ij every half hour.

Use. Same as preceding.

MUCUNA. (Dolichus pruriens. Cowhage. Hairs of the pod.)

Dose. Gr. v-x in syrup, treacle, or honey.

Use. In cases of lumbrici.

OLEUM TEREBINTHINÆ. (See DIURETICS.)

Dose and Form. f ʒ ss-ij, on aromatic water, with hot tincture, (e. g. Tinct. caps. or cinn. or card. c.) ; or as emulsion or linctus.

Use. In cases of tænia, lumbrici.

[**NOTE.**—GEOFFROYA. (G. inermis. The bark.—Dub.)

Dose. Powder ʒ j-3ss.

1. *Decoctum Geoffroyæ.*

Comp. Geoff. ʒ ss, water f ʒ xx, boil to f ʒ x.

Dose. f ʒ ss-ij.]

Use. For lumbrici.

SPIGELIA. (S. Marilandica. Indian pink. The root.)

Dose. Powder gr. x-xx, ʒj-ij, night and morning. Infusion (ʒ ss in Oj.) f ʒ ss-j, f ʒ iv-vj.

Use. For lumbrici and general use.

STANNUM. (Common tin.)

Dose. ʒj-ij- ʒ j, in treacle.

Use. For lumbrici.

From other Classes.

ANTACIDS.

Carb. sodæ.

Liquor calcis.

Liquor potassæ.

PURGATIVES.

Aloes.

Calomel.

Cambogia.

Colocynthis.

Jalapa.

Scammonium.

Tiglia oleum.

TONICS.

Bitters.

Steel.

Antispasmodics.

ÆTHER SULPHURICUS. (See STIMULANTS.)

ACIDUM HYDROCYANICUM. (See NARCOTICS.)

AMMONIACUM. (Gum resin of Dorema A.)
Form and Dose. Gr. v-x-3ss, in pill or emulsion.
 (EXPECTORANTS.)

1. **MISTURA AMMONIACI.** (Lac ammoniaci.)

Comp. Ammoniacum 3ij, water Oss.

Dose. f 3 ss-j.

2. **EMPLASTRUM AMMONIACI.** (See PART II.)

3. **EMPLASTRUM AMMONIACI CUM HYDRARGYRO.**
 (See PART II.)

ASSAFÆTIDA. (Gum resin of Ferula A.)

Dose and Form. Gr. v-3j. in pill, or emulsion ;
 3 ij in 3 xx of water, as enema.

Use. In hysteria, tympanitic distension, tetanus.
 Externally in pertussis, as a plaster.

1. **MISTURA ASSAFÆTIDÆ.**

Comp. Assafætida 3 ij, water Oss.

Dose. f 3 ss-ij (or as *enema* for children.)

2. **TINCTURA ASSAFÆTIDA.**

Comp. Ass. 3 iv, alc. Oij.

Dose. 3 ss-ij.

3. **PILULÆ GALBANI COMPOSITÆ.** (GALBANUM.)

Comp. Assafætida 3 ss, galbanum 3 iss, myrrh
 3 iss, syrup, q. s.

Dose. Gr. x-3j.

4. **SPIRITUS AMMONIÆ FÆTIDUS.**

Comp. Spiritus ammoniæ and assafætida.

Dose. f 3 ss-j (and extemporaneous with sp.
 ammon. and tinct. ass.)

BENZOINUM. (Styrax B. The Balsam.) (EX-
 PECTORANTS. STIMULANTS.)

Dose. Gr. x. (Rarely used.)

1. TINCTURA BENZOINI COMPOSITA.

Comp. Benzoin $\frac{3}{2}$ iiij, storax $\frac{3}{2}$ ij, tolu $\frac{3}{2}$ j, aloes $\frac{3}{2}$ ss, alc. Oij.

Dose. f $\frac{3}{2}$ ss-ij. (Externally—*Friar's Balsam.*)

2. ACIDUM BENZOICUM.

Dose. Gr. v-xx, in powder with gum or sugar.

CAJAPUTI. (See STIMULANTS.)

CAMPHORA. (See NARCOTICS.)

CARDAMINE. (C. Pratensis. The Cuckoo-flower.)

Dose. $\frac{3}{2}$ j-iij of the dried flowers. (Rarely used.)

CASTOREUM. (Preputial secretion of the beaver.)

Form and Dose. Gr. v-xx, in pill, powder, or mixture.

1. TINCTURA CASTOREI.

Comp. Castor $\frac{3}{2}$ ij, alcohol Oij.

Dose. f $\frac{3}{2}$ ss-ij.

CROCUS. (C. Sativus. Saffron. The dried stigmata.)

Dose and Form. Gr. x- $\frac{3}{2}$ ss, in pill or powder. (Ineffective.)

1. SYRUPUS CROCI.

Comp. Saffron $\frac{3}{2}$ v, sugar $\frac{3}{2}$ xvij, water f $\frac{3}{2}$ x.

Dose. f $\frac{3}{2}$ j-ij, chiefly for colouring.

CUPRI AMMONIO-SULPHAS. (See TONICS.)

GALBANUM. (Gum resin of G. Officinale.)

Dose and Form. Gr. v-x- $\frac{3}{2}$ ss, in pill or emulsion.

1. PILULÆ GALBANI COMPOSITÆ.

Comp. Assafætida $\frac{3}{2}$ ss, galbanum $\frac{3}{2}$ iss, myrrh,
 $\frac{3}{2}$ iss, syrup q. s.

Dose. Gr. x- $\frac{1}{2}$ j.

2. EMPLASTRUM GALBANI. See PART II.

MASTICHE. (Pistacia Lentiscus. The resin.
 Gum Mastic.) (DIURETICS. EXPECTORANTS.)

Dose. Gr. v-x. [Also to divide more active
 medicines, as aloes in the "Pilulæ ante cibum"
 of the Parisian Codex.] (See ALOES EXT.)

MOSCHUS. (Preputial secretion of the musk
 animal.)

Dose and Form. Gr. v-xv, in pill, bolus, or emul-
 sion.

Use. Hiccough, epilepsy, asthma, pertussis, palpi-
 tation.

1. MISTURA MOSCHI.

Comp. Musk, gum, sugar, $\ddot{\text{aa}}$ $\frac{3}{2}$ iij, rose water,
 $f\frac{3}{2}$ xx.

Dose. $f\frac{3}{2}$ j-iss, ($f\frac{3}{2}$ j contains gr. ix.)

[MOSCHUS FACTITIUS. (Artificial Musk.)

Comp. Acid. nit. fort. $\frac{3}{2}$ iiiss, ol. Succini $\frac{3}{2}$ j.

Form. Tincture: Mosch. fact. $\frac{3}{2}$ j, spir.
 rect. $\frac{3}{2}$ x.

Dose. $\frac{3}{2}$ j.]

OLIBANUM? (Gum resin of Boswellia thuri-
 fera. Frankincense.)

Dose. Gr. x- $\frac{3}{2}$ ss, in emulsion, with yolk of egg.

OPOPOONAX. (Gum res. of O. Chironium.)

Dose. Gr. x- $\frac{3}{2}$ ss, in emulsion. (Rarely used.)

PETROLEUM? (Barbadoes tar.) (EXPECTO-
 RANTS.)

Dose. ℥x-3 ss, on sugar, in emulsion, or with aromatic waters.

SAGAPENUM. (Gum resin of Ferula S.)

Dose and Form. Gr. v-3j, in pill.

1. PILULÆ GALBANI COMPOSITÆ. (See GALBANUM.)

2. PILULÆ SAGAPENI COMPOSITÆ.

Comp. 16 sagapenum, 1 aloes, syrup of ginger, q. s.

Dose. Gr. v-x-xx (17 grs. contain 1 of aloes.)

STYRAX. (See EXPECTORANTS.)

SUCCINUM. (AMBER.)

1. OLEUM SUCCINI.

Dose. ℥v-x in mucilaginous fluids.

VALERIANA. (Root of V. officinalis.)

Dose. 3j-3 ss, in powder.

Use. Hysteria, hemicrania, pervigilium, low fevers.

1. TINCTURA VALERIANÆ.

Comp. Valerian 3 iv, dil. alc. Oij.

Dose. f 3 ss-ij.

2. TINCTURA VALERIANÆ COMPOSITA.

Comp. Valerian 3 iv, aromat. sp. of ammonia, Oij.

Dose. f 3 j-ij. in sweetened water or milk.

Use. In Hysteria.

ZINCI OXIDUM. (See TONICS.)

Aromatics. Aromatic Stimulants.

CANELLA. (C. alba. Wild Cinnamon. The Bark.)

Form and Dose. Powder gr. x-3 ss—used chiefly in the preparation of other formulæ.

CAPSICUM. (C. annum. Cayenne Pepper.

The berry.)

Form and Dose. Powder (in pills with bread crumb) gr. v-x; infusion (caps. 3 ij, water f $\frac{2}{3}$ x,) f $\frac{2}{3}$ ss-j; gargle, with water or vinegar. **Use.** In malignant sore throat.

1. TINCTURA CAPSICI.

Comp. Caps. 3 i, dil. alc. Oij.

Dose. f 3 j-3 ij. As gargle 3 j, infus. rosæ 3 vj.

Use. In malignant scarlatina, typhus, locally to relaxed uvula.

CARDAMOMUM. (Alpinia C. The seeds.)

Form and Dose. Gr. v- $\frac{3}{4}$ j in powder. (Rarely used alone, but much in composition.)

1. TINCTURA CARDAMOMI.

Comp. Card. 3 iv, dil. alc. Oij.

Dose. f 3 j-ij, chiefly as an adjunct.

2. TINCTURA CARDAMOMI COMPOSITA.

Comp. Card. and carui seeds, of each 3 iiiss, cochineal 3 j, cinnamon 3 v, raisins 3 v, proof sp. f $\frac{2}{3}$ xl.

Dose. 3 j-ij.

CARYOPHYLLUS. (C. Aromaticus. The Clove Tree, The unripe flowers.)

Dose. Gr. v-x in powder. (Little used.)

Use. To allay nausea. To correct flatulence. To excite languid digestion.

1. INFUSUM CARYOPHILLI.

Comp. Cloves 3 ii, water Oi.

Dose. 3 j-ij. Combined with other med.

Incomp. Salts of iron and zinc, lime water.

2. OLEUM CARYOPHILLI.

Dose. $\frac{1}{2}$ j-ij, with pills.

CINNAMOMUM. (Laurus. C. Cinnamon. The bark.)

Dose. Gr. x- $\frac{3}{2}$ ss in powder.

Use. To allay nausea, to remove flatulence.

1. PULVIS CINNAMOMI COMPOSITUS.

Comp. 2 cinnamon, 1 cardamoms, 2 ginger, nutmeg grated 1 part.

Dose. Gr. x- $\frac{3}{2}$ ss. As a corrigent.

2. AQUA CINNAMOMI.

Comp. From the oil, or distilled from the bark.

Dose. Ad libitum, as a vehicle.

3. TINCTURA CINNAMOMI.

Comp. Cinn. $\frac{2}{3}$ iii, dil. alc. Oij.

Dose. f $\frac{3}{2}$ j- $\frac{2}{3}$ ss.

4. TINCTURA CINNAMOMI COMPOSITA.

Comp. Cinn. $\frac{2}{3}$ j, cardam. $\frac{2}{3}$ ss, ginger 3 ijj, dil. alc. Oij.

Dose. f $\frac{3}{2}$ j-ij. Used as the last.

5. OLEUM CINNAMOMI.

Dose. $\frac{1}{2}$ ij-ijj, as an adjuvant and corrigent.

6. SPIRITUS CINNAMOMI.

Comp. Oil of cinn. 3 ij, proof spir. 1 gal., water f $\frac{2}{3}$ xx, distil to 1 gall.

Dose. 3 j-ijj.

CONFECTIO AROMATICA.

Comp. Pulv. aromat. $\frac{2}{3}$ vss, saffron $\frac{2}{3}$ ss, syr. aurant $\frac{2}{3}$ vj, mel. desp. $\frac{2}{3}$ ij.

Dose. Grs. x- $\frac{3}{2}$ j,

(Pulv. aromat. is composed cinnam. and ginger, each $\frac{2}{3}$ ij, cardam. and nutmeg each $\frac{2}{3}$ j.

CONTRAJERVA. (Dorstenia C. The root.)

Form and Dose. Powder 3j-3ss; infus. (Contr. 3ij, water 3vj) 3j-ij. (Rarely used.)

CUBEBA. (Piper Cubeba. The berries.)

Dose. Powdered—Gr. v-x-3ij.

Use. As a stomachic in disorders of digestive organs.

1. **TINCTURA CUBEBAE.**

Comp. Cubebs 3iv, dil. alc. Oij.

Dose. f3j-ii.

Use. In advanced stage of gonorrhœa.

INULA. (I. Helenum. Elecampane. The root.)

Form and Dose. Powder 3j-3ij; decoction (3ss, water f3xx,) f3j-ij.

MYRISTICA. (Nutmeg. The nut of the Myristica Moschata, and the Arillus, Mace.)

Dose. Gr. x-3ss.

1. **OLEUM MYRISTICÆ.** (See PART II.)

Dose. Mij-vj. (Externally stimulant.)

2. **SPIRITUS MYRISTICÆ.**

Comp. Nutmegs 3ii, dil. alc. 1 gall., water Oj, distil to 1 gall.

Dose. f3j-3ij.

PIMENTA. (Myrtus P. Allspice. The unripe berries.)

Dose. Powder, gr. x-3j.

Use. In cases of flatulence.

1. **OLEUM PIMENTÆ.**

Dose. Mij-vj.

2. **SPIRITUS PIMENTÆ.**

Comp. Pimenta 3ii, dil. alc. 1 gall., water Oj, distil to 1 gall.

Dose. f 3ij-ij.

3. AQUA PIMENTÆ.

Comp. Pimenta ℥vj, or ol. pimenta, 3ij, dil. alc. f 3 vij, water 2 gall. distil to 1 gall.

Dose. f 3 j-ij. (A vehicle and flavourer.)

PIPER LONGUM. (Long Pepper. The unripe spadix.)

Dose. Gr. v-3j. (A condiment.)

PIPER NIGRUM. (Black Pepper.)

Dose. Gr. v-3j. In powder or pills.

Use. In flatulence or debility of stomach.

1. CONFECTIO PIPERIS NIGRI. (Ward's Paste.)

Comp. Pip. nig., inula (elecampane) aa 3 iv. fænic. sem. 3 xij, mell. et sacch. aa 3 viij.

Use. In piles unattended with inflammation.

SINAPIS. (S. nigra. The seed.) (EMETICS.)

Form and Dose. Mustard whey (bruised seeds or powder 3 ss, milk 3 xx-strain) 3 iv.

1. CATALPLASMA SINAPIS. (See PART II.)

ZINZIBER. (The rizoma or root.)

Dose. Powder gr. v-x-3j; infusion (ginger tea—Ginger 3 ij-3 ss, water f 3 vj), f 3 ss-j.

Use. In dyspepsia. The state of alimentary canal attending atonic gout.

1. TINCTURA ZINZIBERIS.

Comp. Ginger 3 viij, alc. Oij.

Dose. 3 j-ij.

2. SYRUPUS ZINZIBERIS.

Comp. Tinc. Zinzib. f 3 iv, syrp. Cong. j.

[EXTRACTUM ZINZIBERIS. (Non-officinal.)

Dose. Gr. ss-iiij.]

*Aromatic Bitters. Stimulant Bitters.***ABSINTHIUM.** (Artemisia A. Wormwood.

The flowering summits.)

Dose and Form, Powder $\frac{1}{2}$ j- $\frac{3}{2}$ j; infusion ($\frac{2}{3}$ j
in $\frac{2}{3}$ xx water) $\frac{2}{3}$ ss-ij; extract (Dub.) gr. x.- $\frac{1}{2}$ j.**Incomp.** Sulphates of iron and zinc.[**OLEUM ABSINTHII.** (Not officinal.)**Dose.** $\frac{1}{2}$ j-v.]**Use.** In debilitated states of the digestive organs.**ACORUS.** (A. calamus. Sweet Flag. The root.)**Dose and Form.** Powder $\frac{1}{2}$ j- $\frac{3}{2}$ j; infusion (Acor.
 $\frac{2}{3}$ j, water f $\frac{2}{3}$ xx) f $\frac{2}{3}$ j-ij, joined with other tonics.**Use.** In uneasiness from flatulence.**ANTHEMIS.** (A. nobilis. Chamomile flowers.)**Dose and Form.** Powder, gr. x- $\frac{3}{2}$ ss, (seldom used); extract (Ed. Dub. watery) gr. v-x- $\frac{1}{2}$ j.**Use.** Remission of fever.1. **INFUSUM ANTHEMIDIS.****Comp.** Chamomile $\frac{2}{3}$ ss, boiling water Oj.**Dose.** f $\frac{2}{3}$ j-ij.**Incomp.** Iron, bichlor. hydr., infus. cinch., isinglass.**Use.** In enfeebled digestion; general debility.2. **OLEUM ANTHEMIDIS.****Dose.** $\frac{1}{2}$ j-v. (Stimulant. Antispasmodic.)**AURANTII CORTEX.** (Citrus vulgaris. Bitter Orange. The peel.)**Dose.** Powder gr. x- $\frac{3}{2}$ ss. (Rarely used.)

1. INFUSUM AURANTII COMPOSITUM.

Comp. Orange peel $\frac{3}{2}$ ss, lemon peel (fresh) $\frac{3}{2}$ ij;
cloves $\frac{3}{2}$ j, water f $\frac{3}{2}$ xx.

Dose. f $\frac{3}{2}$ j-ij.

Incomp. Sulph. ferr., infus. cinch. cord., lime
water.

2. TINCTURA AURANTII.

Comp. Orange peel $\frac{3}{2}$ iiiss, proof spirit f $\frac{3}{2}$ xl.

Dose. f $\frac{3}{2}$ j-ij.

3. CONFECTIO AURANTII.

Comp. Orange peel $\frac{1}{2}$ bj, sugar $\frac{1}{2}$ bij.

Dose. $\frac{3}{2}$ j-ss, and as a vehicle for powders; e. g.
ferr. sesquioxyd, &c.

4. SYRUPUS AURANTII.

Comp. Or. peel $\frac{3}{2}$ ii, sugar $\frac{1}{2}$ bijss. boiling water
Oj.

Dose. $\frac{3}{2}$ j-ij. Chiefly to flavour saline and sto-
machic draughts.

5. AQUA FLORUM AURANTII.

Comp. Or. flowers $\frac{3}{2}$ xxx, spir. rect. f $\frac{3}{2}$ xiv, water
 $\frac{1}{2}$ gall. For flavour, and as a vehicle only.

6. OLEUM AURANTII.

Dose. M v-x. Stimulant. For flavour.

AURANTIUM. (Citrus A. Sweet orange.
The peel.)

Form and Dose. Powder gr. x- $\frac{3}{2}$ ss. Infusion as
a substitute for the bitter orange.

CASCARILLA. (Croton C. The bark.)

Dose. Powder gr. x- $\frac{3}{2}$ ss.

Use. In flatulent colic, dyspepsia, chronic dia-
rhœa.

1. INFUSUM CASCARILLÆ.

Comp. Casc. $\frac{2}{3}$ i, water Oj.*Dose.* f $\frac{2}{3}$ j-ij.

2. TINCTURA CASCARILLÆ.

Comp. Casc. $\frac{2}{3}$ iiiss, proof sp. f $\frac{2}{3}$ xx.*Dose.* f $\frac{2}{3}$ j-ij.

3. MISTURA CASCARILLÆ COMPOSITÆ.

Comp. Infus. casc. (No. 1) f $\frac{2}{3}$ viiiss, tinct. camph.
c. f $\frac{2}{3}$ j, acet. scillæ f $\frac{2}{3}$ ss.*Dose.* f $\frac{2}{3}$ j-iss. (EXPECTORANTS.)

CUSPARIA. (Angustura Bark.)

Dose. Powder gr. x- $\frac{3}{2}$ ss. (Rarely used.)

1. INFUSUM CUSPARIÆ.

Comp. Cusp. $\frac{3}{2}$ v, water f $\frac{2}{3}$ xx.*Dose.* $\frac{2}{3}$ j-ij (with tinct. cinn. added).

LIMONES. (Citrus Limonum. Lemon. The fruit.)

1. LIMONUM CORTEX. (Lemon peel.)

Form and Dose. Powder gr. x- $\frac{3}{2}$ j. (Rarely used except in other preparations, q. v.)

2. OLEUM LIMONUM.

Dose. $\frac{1}{2}$ j-v.

LUPULUS. (Humulus L. The Hop. The dried strobiles.)

Form. In substance as a pillow. The Lupulinic grains or yellow powder. And—

1. INFUSUM LUPULI.

Comp. Hops $\frac{2}{3}$ ss, water Oj.*Dose.* f $\frac{2}{3}$ j-ij.*Incomp.* Acids, alkalies, sulph. ferr., tart. emet., alcohol.

2. TINCTURA LUPULI.

Comp. Hops $\frac{3}{2}$ v, dil. alc. Oij.

Dose. f $\frac{3}{2}$ ss-ij.

3. EXTRACTUM LUPULLI. (A watery extract.)

Dose. Gr. v- $\frac{3}{2}$ j.

[NOTE. *Lupulina.* Yellow powder. (Not officinal.)]

Dose and Form. Powder or pills gr. vii-xij.

a. *Tinctura Lupulinæ.*

Comp. Lupulinic grains $\frac{3}{2}$ iv, spirit Oij.

Dose. f $\frac{3}{2}$ ss-ij.]

Use. In delirium and wakefulness of drunkards.

Astringents.

ACIDUM SULPHURICUM DILUTUM.

(See TONICS, REFRIGERANTS.)

Use. In colliquative sweats, passive hemorrhage, diarrhoea.

ACIDUM ACETICUM. (REFRIGERANTS.)

ALUMEN. (Alum, sulphate of alumina and potassa.)

Dose. Gr. v- $\frac{3}{2}$ j.

Form. Solution (in aromatic water: an oz. of water dissolves $\frac{3}{2}$ ss), pill, powder.

[NOTE. Alum whey: boil powdered alum $\frac{3}{2}$ ij in a pint of milk, and strain.

Dose. A wine-glassful.]

Incomp. Alkalies and their carbonates, tartrate of potash, vegetable astringents.

Use. In hemorrhages ; as a gargle, as a collyrium.

1. ALUMEN EXSICCATUM. (Dried alum.)

Dose. Gr. v-x.

Form. Solution, pill.

Use. As an escharotic to fungous ulcers.

2. LIQUOR ALUMINIS COMPOSITUS.

(See PART II.)

ARGENTI NITRAS. (See TONICS.)

BORAX. (Sodæ Biboras.)

Dose. 3j- 3 ss.

(Rarely used internally.)

1. MEL BORACIS.

Comp. Borax 3 j, honey f 3 j.

Form. As a gargle, diluted,

Use. For aphthæ of children.

CATECHU. (Acacia Catechu. Extract of the wood.)

Dose. Gr. x- 3j.

Form. Powder, mixture (with syrup and gum), bolus.

Use. Diarrhœa, uterine hemorrhage, relaxed uvula, spongy gums, indolent ulcers, leucorrhœa, gonorrhœa, epistaxis.

Incomp. Alkaline salts destroy its astringency.

1. INFUSUM CATECHU COMPOSITUM.

Comp. Catechu 3 ss, cinnamon 3j, water Oj.

Dose. f 3 j-ij.

2. TINCTURA CATECHU.

Comp. Catechu 3 iii, cinnamon 3 ii, dil. ale. Oij.

Dose. f 3 j-ij in chalk mixture, or other liquids.

CINCHONA. (See TONICS.)**CUPRUM.****1. CUPRI SULPHAS.**

Dose and Form. Gr. $\frac{1}{8}$ - $\frac{1}{4}$ -ss-ij (combined with opium.)

Use. In chronic diarrhoea, ill-conditioned ulcers, collyrium in chronic ophthalmia.

FERRUM. (See TONICS.)

Use. In debility from inordinate discharges.

1. FERRI SULPHAS.**2. MISTURA FERRI COMPOSITA.****3. PILULÆ FERRI COMPOSITÆ.****4. TINCTURA FERRI SESQUICHLORIDI.**

Use. In gleet, passive hemorrhages.

GALLÆ. (Oak Galls. Gemmæ Morbidæ.)

Form and Dose. Powder gr. x-xx. Infusion (galls f $\frac{2}{3}$ ss, boiling water f $\frac{2}{3}$ vj) f $\frac{2}{3}$ ss-iss.

Use. In chronic diarrhoea, as a gargle in relaxed uvula.

1. TINCTURA GALLÆ.

Comp. Galls $\frac{2}{3}$ iv, dil. alc. Oij.

Dose. f $\frac{2}{3}$ ss-ij in fluids, as gargle; or lotion, diluted with water.

Incomp. Metallic salts, cinchona.

2. UNGUENTUM GALLÆ COMPOSITUM. (See PART II.)

Use. In hemorrhoids.

GRANATUM. (*Punica G.* Pomegranate. Rind of the fruit.)

Dose and Form. Powder $\frac{2}{3}$ ss-j.

1. DECOCTUM GRANATI.

Comp. Granat. $\frac{2}{3}$ ij, water f $\frac{2}{3}$ xxx (to f $\frac{2}{3}$ xx.)

Dose. f $\frac{2}{3}$ iss-ij.

(Bark of the Root ANTHELMINTIC, q. v.)

HÆMATOXYLUM. (H. Campechianum. Log-wood.)

1. EXTRACTUM HÆMATOXYLI.

Dose. Gr. x- $\frac{1}{3}$ ss.

Form. Pill, mixture, with carminative waters.

[DECOCTUM HÆMATOXYLI—Edinb.

Comp. Logwood $\frac{2}{3}$ i, water Oij.

Dose. f $\frac{2}{3}$ j-ij. (For children f $\frac{2}{3}$ ij- $\frac{2}{3}$ ss.)]

Use. In relaxed condition succeeding cholera infantum.

KINO. (Extract of *Pterocarpus erinaceus*.)

Dose and Form. Powder, gr. v-x-xx. Mixture with mucilage and the aromatic waters.

Incomp. Acids, alkalies, sulphate of iron.

1. TINCTURA KINO.

Comp. Kino $\frac{2}{3}$ iiiss, rectified spirit f $\frac{2}{3}$ xl.

Dose. f $\frac{2}{3}$ j-ij, with chalk mixture.

Use. In diarrhoea without fever.

2. PULVIS KINO COMPOSITUS.

Comp. 15 kino, 4 cinnamon, 1 opium, (1 gr. of opium in $\frac{1}{2}$ j.)

Dose. Gr. v- $\frac{1}{2}$ j, in mixture.

KRAMERIA. (K. triandria. Rhatany root.)

Dose. Gr. x- $\frac{1}{3}$ ss in powder, mixture, or electuary.

Use. Chronic diarrhoea, leucorrhœa, passive hemorrhage.

1. INFUSUM KRAMERIÆ.

Comp. Krameria $\frac{2}{3}$ j, water Oj.

Dose. f $\frac{3}{4}$ j-ij.

Incomp. Sulphate of iron.

[**EXTRACTUM KRAMERIAE** (Ed. Dub.)

Dose. Gr. x-xx, pill, mixture.]

OPIUM. (See **NARCOTICS.**)

PLUMBUM.

1. PLUMBI ACETAS.

Dose. Gr. ss-iv.

Form. Pill with opium, solution with vinegar and distilled water.

Incomp. Sulphuric acid, sulphates, liq. ammoniae acetatis, &c.

Use. In hemorrhages, dysentery, diarrhoea.

QUERCUS. (Q. pedunculata. The oak. The bark.)

Dose. Powder $\frac{3}{4}$ ss-ij.

Use. Diarrhoea, leucorrhœa.

1. DECOCTUM QUERCUS.

Comp. Oak bark $\frac{3}{4}$ j, water Ojss (to Oj.)

Dose. f $\frac{3}{4}$ ij-iv. (Externally as gargle, injection, lotion.)

ROSA. (Rosa Gallica. The red rose petals.)

1. INFUSUM ROSÆ COMPOSITUM.

Comp. Rose $\frac{3}{4}$ ss, water Oijss, sugar $\frac{3}{4}$ iss, dilute sulphuric acid $\frac{3}{4}$ ij.

Dose. $\frac{3}{4}$ j-ij. (Externally as gargle.)

Incomp. Alkalies and earths, sulphate of iron.

Use. In hemorrhages and colliquative sweats.

2. CONFECTIO ROSÆ GALLICÆ.

Comp. Rose $\frac{3}{4}$ iv. sugar $\frac{3}{4}$ xxx, mel. $\frac{3}{4}$ vj, aq. ros. $\frac{3}{4}$ viij.

Dose. $\frac{3}{4}$ j-ij, as a vehicle for calomel, quinine, &c.

3. MEL ROSÆ.

Comp. Rose $\frac{3}{4}$ ij, water Oiss, honey Oij. (A honeyed decoction of Rose.)

Dose. $\frac{3}{4}$ j-iv. (Chiefly in gargles.)

[**TINCTURA ROSÆ.** An admirable substitute for the infusion is prepared under this name. It is made by digesting 1 oz. of red rose petals with 4 oz. of proof spirit made with rose-water. Of this tincture $\frac{M}{M}$ xxx, with $\frac{M}{M}$ xij of acid. nit. dil., is equal to $\frac{3}{4}$ iss of infusion of roses.

TORMENTILLA. (Tormentil or Septfoil. The root.)

Dose and Form. Powder $\frac{3}{4}$ ss-j.

1. DECOCTUM TORMENTILLÆ.

Comp. Tormentil $\frac{3}{4}$ ij, water f $\frac{3}{4}$ xxx (to f $\frac{3}{4}$ xx.)

Dose. f $\frac{3}{4}$ j-ij (used also as an injection in leucorrhœa.)

ULMUS. (U. campestris. The bark.)

1. DECOCTUM ULMI.

Comp. Fresh elm bark $\frac{3}{4}$ iiss, water $\frac{3}{4}$ xl (to xx.)

Dose. $\frac{3}{4}$ ij-iv. (Rarely used.)

UVA URSI. (Leaves of the bear-berry.)

Dose. $\frac{1}{2}$ j- $\frac{3}{4}$ j of the powder (too bulky).

Use. In diabetes, catarrh of bladder, incontinence of urine.

1. DECOCTUM UVÆ URSI.

Comp. Uva ursi $\frac{3}{4}$ j, water f $\frac{3}{4}$ xx (to f $\frac{3}{4}$ xvi.)

Dose. f $\frac{3}{4}$ j-ij.

2. EXTRACTUM UVÆ URSI.

Dose. Gr. v-xv.

ZINCUM.

1. ACETAS ZINCI.

Dose. Gr. j-ij. (Rarely used internally.)

2. SULPHAS ZINCI.

Dose. Gr. j-v. (Much used externally.)*Use.* Leucorrhœa, gonorrhœa, ophthalmia, and indolent ulcers.*Carminatives.**Use.* As a class, to prevent nausea, to correct flatulence, and as corrigents to other medicines.

ANETHUM. (A. Graveolens. Dill. The fruit.)

Dose. Powder gr. v- 3 ss.

1. AQUA ANETHI.

Comp. Anethum 3 xvij, proof sp. f 3 vij, water 2 gall. (or by diffusing the oil in water.)*Dose.* f 3 j-ij (infants)-f 3 j-ij (adults.)

ANISUM. (Pimpinella A. Anise. The seed.)

Dose. Powder gr. x- 3 ss.

1. OLEUM ANISI.

Dose. M v-xv. (on sugar, or with sugar in mixture.)

2. SPIRITUS ANISI.

Comp. Anise 3 x, proof sp. 1 gall., water f 3 xl.*Dose.* f 3j- 3 ss.

CARUI. (Carum C. Caraway. The seeds.)

Dose. Powder gr. x-3ss.

1. OLEUM CARUI.

Dose. ℥ij-v.

2. SPIRITUS CARUI.

Comp. Carui ʒ xj, proof spir. ½ gall., water f ʒ xx,
distil to ½ gall.

Dose. f ʒ j-ʒ ss.

3. AQUA CARUI.

Comp. Carui ʒ xvij, water 2 gall., spir. f ʒ vij,
distil to 1 gall.

Dose. f ʒ j-iv.

CORIANDRUM. (C. Sativum. Coriander. The
seeds.)

Dose. Powder gr. x-3j-3j.

CYMINUM. (Cuminum Cyminum. Cumin.
The seeds.)

Dose. Powder gr. x-3ss. (Rarely used.)

DAUCUS. (D. Carota. Carrot. The seed.)

Dose. Powder gr. x-3j-3 ss.

FENICULUM. (F. Vulgare. Fennel. The
seeds.)

Dose. Gr. x-3 ss.

1. AQUÆ FENICULI.

Dose. f ʒ j-iij.

LAVANDULA. (Flower tops of L. vera. Lav-
ender.)

Dose. Powder 3j-3 ss. (Rarely used.)

1. OLEUM LAVANDULÆ.

Dose. ℥ij-v.

2. SPIRITUS LAVANDULÆ.

Comp. Lavender $\frac{1}{2}$ ij, alc. 1 gall, water Oij.

Dose. f 3 ss-ij. (Rarely used.)

3. TINCTURA LAVANDULÆ COMPOSITA. (Red Lavender.)

Comp. Spir. lavand. Oijj, spir. of rosemary Oj,
cinnamon $\frac{1}{2}$ j, nutmeg $\frac{1}{2}$ ss, red saunders 3 ij,
cloves $\frac{1}{2}$ ij.

Dose. f 3 ss-j. (On a lump of sugar.)

LAURUS. (L. Nobilis. Sweet Bay. The berries and leaves.)

Dose. Of the berries powdered 3 ss-j. The oil.
Infusion. (Rarely used.)

MENTHA PIPERITA. (Peppermint.)

Dose. Powder gr. x- 3 ss; infusion (3 ij water
f $\frac{1}{2}$ viij) f $\frac{1}{2}$ j-ij. (Rarely used.)

1. OLEUM MENTHÆ PIPERITÆ.

Dose. $\frac{1}{2}$ ij-v.

2. SPIRITUS MENTHÆ PIPERITA.

Comp. Ol. menthæ 3 ij, proof sp. 1 gall., water
f $\frac{1}{2}$ xx.

Dose. f 3 ss-ij.

[NOTE.—Essence of peppermint is made by dissolving 3 j of the oil in $\frac{1}{2}$ j of rectified spirit.]

3. AQUA MENTHÆ PIPERITÆ.

Comp. Ol. menth. pip. 3 ss, carb. magnes. 3 ss,
aq. dist. Oij.

Dose. f $\frac{1}{2}$ j-ij.

MENTHA PULEGIUM. (Pennyroyal.)

Dose. Powder (rarely used), 3 ss-j.

1. OLEUM MENTHÆ PULEGIL.

Dose. ℥ij-v.

2. SPIRITUS MENTHÆ PULEGII.

Comp. Ol. Puleg. ʒ iiij, proof spir. 1 gall., water f ʒ xx.*Dose.* f ʒ ss-ij.

3. AQUA MENTHÆ PULEGII.

Comp. Pennyroyal (dried) ʒ xij (or oil ʒ j), proof spir. f ʒ iiiss, water 1 gall.

MENTHA VIRIDIS. (Spearmint.)

Dose and Form. Powder ʒ ss-ij (rarely used); infusion (ʒ ij-water f ʒ viij), f ʒ j-ij.

1. OL. MENTHÆ VIRIDIS.

Dose. ℥ij-v.

2. SPIRITUS MENTHÆ VIRIDIS.

Comp. Oil ʒ iiij, spir. 1 gall., water f ʒ xx.*Dose.* f ʒ ss-ij.

3. AQUA MENTHÆ VIRIDIS.

Comp. See Aq. Menth. Pip.*Dose.* f ʒ j-ij.

ORIGANUM. (Marjoram.)

Dose. Powder, gr. x-ʒ ss; infusion (ʒ ij, water f ʒ viij) f ʒ j-ij. (Rarely used.)

1. OLEUM ORIGANI.

Dose. ℥ij-v.

ROSMARINUS. (Rosemary.)

Dose. Powder, gr. x-ʒ ss (rarely used); infusion (ʒ ij, water f ʒ viij) f ʒ j-ij.

1. OLEUM ROSMARINI.

Dose. ℥ij-v.

2. SPIRITUS ROSMARINI.

Comp. Oil 3 ij, alc. 1 gall., water Oj.*Dose.* f 3 ss-ij.*Diaphoretics.*AMMONIÆ SESQUICARBONAS ? (See RE-
FRIGERANTS.)*Dose.* Gr. iij-vj.

1. LIQUOR AMMONIÆ ACETATIS.

Comp. Sesq. am. 3 ix, (vel q. s.) Acet. dest.
f 3 xx. A saturated solution.*Dose.* f 3 iij-vj- 3 j.*Incomp.* Acids, potass, sugar.*[Saline Draughts.]* Ammon. sesquicarb. 3 j, acid.
citric. gr. xxiv, *vel* acid. tart. gr. xxv, *vel* succ.
lim. 3 vj.]*Use.* In febrile and inflammatory diseases.

ANTIMONIUM. (See EMETICS.)

1. ANTIMONII POTASSIO-TARTRAS.

Dose. Gr. 1½-ij.

2. VINUM ANTIMONII POTASSIO-TARTRATIS.

Comp. Ant. pot. tart. 3 j, sherry f 3 x (gr. j, in
f 3 ss).*Dose.* M x-xv- 3 j.3. PULVIS ANTIMONII COMPOSITUS. (Pulvis Anti-
monialis.)*Comp.* 2 antimonious acid, 3 phosph. of lime.*Dose.* Gr. iij-x.*Form.* Pill, powder.

DIOSMA. (D. crenata. Buchu. The leaves.) ?
Dose. (Of the powder) 3j-3ss.

1. INFUSUM DIOSMÆ.

Comp. Diosma 3j, water Oj.

Dose. f 3j-ij.

DULCAMARA? (Solanum D. The stalk.)

1. DECOCTUM DULCAMARÆ.

Comp. Dulc. 3j, water Oiss-Oj.

Dose. f 3ss-iss.

GUAIACUM. (With Diluents.)

1. THE RESIN.

Dose. Gr. v-xx.

Form. Pill, bolus, mixture.

Use. In rheumatism.

2. MISTURA GUIACI.

Comp. Guaiacum 3 iij, sugar 3 ss, mucilage
3 ss, cinnamon water, f 3 xix.

Dose. f 3ss-ij (guaiac. gr. ix in f 3j).

3. TINCTURA GUIACI.

Comp. Guaicum lb ss, alc. Oij.

Dose and Form. f 3j-ij, in mucilage, or syrup
and water.

Incomp. Water (decomposes it).

3. TINCTURA GUIACI COMPOSITA.

Comp. Guaiacum 3 iv, aromatic spirit of am-
monia Ojss.

Dose. f 3ss-ij in mucilage, or syrup and water.

Use. Chronic rheumatism, amenorrhœa.

IPECACUANHA. (Cephaelis I. The root.)

1. PULVIS IPECACUANHÆ. The powdered root.

Dose. Gr. ss-j.

2. PULVIS IPECACUANHÆ COMPOSITUS.

Comp. 1 ipecacuan, 1 opium, 8 sulphate of potass, (gr. x, contain 1 gr. of opium).

Use. In cases where not much fever, cerebral disease, or sick stomach.

Dose. Gr. iiiss-v-xx.

Form. Powder, pill, mixture.

3. VINUM IPECACUANHÆ.

Comp. Ipecacuanha ȝ ij, sherry Oij.

Dose. ȝ xx-ȝ j.

Use. For children.

4. PILULÆ IPECACUANHÆ COMPOSITÆ.

Comp. 3 Pulv. ipec. comp., 1 squill, 1 ammoniac, mucilage q. s.

Dose. Gr. v-x ($\frac{1}{2}$ gr. opium in 10 grs).

MEZEREUM? (Daphne M. Bark of the root.) (STIMULANTS, DIURETICS.)

Form and Dose. Decoction (ȝ j in ȝ xxx water, to ȝ xx) ȝ ij-vj.

Use. In secondary syphilis, cutaneous diseases.

SASSAFRAS? (See ALTERATIVES.)

Dose. Of the powder ȝ ss-j. (Rarely used.)

SENEGA? (See EXPECTORANTS, ALTERANTS.)

Dose. (Of the powder) gr. x-xx. (Rarely used.)

1. DECOCTUM SENEGRÆ.

Comp. Senega root ȝ j, water Oiss-Oj.

Dose. f ȝ j-ij.

Use. In rheumatism.

SULPHUR?

1. SULPHUR PRECIPITATUM.

Dose. 3 ss-j, as an electuary, or suspended in milk.

Use. In chronic rheumatism.

*Diluents. Demulcents. Emollients.
Ptisans.*

ACACIA. (Exudation from *A. vera*. Gum Arabic.)

Dose. Əss-j-ij.

Form. Powder, mixture.

Incomp. Acet. plumb., æther, spirits.

1. **MISTURA ACACIÆ.** (Mucilago acaciæ.)

Comp. Acacia ʒ x, boiling water f ʒ xx.

Dose. f ʒ j- ʒ j.

Incomp. Acet. plumb., æther, tinctures.

ALTHÆA. (*A. officinalis*. Marshmallow. The root.)

Form. Decoction (*A.* ʒ iss. water ʒ xx).

Dose. Of D. ad libitum, ʒ j-ij subinde.

1. **SYRUPUS ALTHÆÆ.**

Comp. Decoction of Althaea with sugar.

Dose. ʒ j-ij. (Readily ferments.)

AMYGDALA AMARA. (Bitter almonds.)

AMYGDALA DULCIS. (Sweet almonds.)

Use. In catarrhal affections.

1. **CONFECTIO AMYGDALÆ.** (Almond confect.)

Comp. Amygd. ʒ viij, acac. pulv. ʒ j, sacch. ʒ iv.

2. **MISTURA AMYGDALÆ.** (Almond emulsion.)

Comp. Sweet almonds $\frac{3}{2}$ ss, pulv. acac. $\frac{3}{2}$ ss,
sugar $\frac{3}{2}$ ij, water $\frac{3}{2}$ viij.

Dose. $\frac{3}{2}$ j-ij, ad libitum.

Use. In catarrhal and dysenteric affections.

3. OLEUM AMYGDALÆ. (Oil of almonds. Ex-
pressed from both kinds.)

Dose and Form. $\frac{3}{2}$ j-ij, in emulsion.

AMYLUM. (Fecula Sem. tritici. Starch.)

Form. Powder (externally); mixed in water.

1. DECOCTUM AMYLI. (Mucilago amyli. Starch
emulsion.)

Comp. Amylum $\frac{3}{2}$ ss, water $\frac{3}{2}$ xx.

Dose. Ad libitum (for enema).

Incomp. Preparations of iodine.

AVENA. (A. sativa. The common oat.)

Form. The meal in decoction, poultice, &c.

CETACEUM. (From the head of the Sperma-
ceti whale. Spermaceti.)

Form and Dose. $\frac{3}{2}$ ss-iss in an emulsion with
egg or sugar.

Use. In catarrh.

CETRARIA. (Lichen Islandicus. Iceland moss.)

1. DECOCTUM CETRARÆ.

Comp. Cetraria $\frac{3}{2}$ ss, water Oiss, boil to Oj.

Dose. f $\frac{3}{2}$ j-iv, ad libitum.

Use. In phthisis.

CYDONIA. (C. vulgaris. Common quince.
The seeds.)

1 DECOCTUM CYDONIÆ.

Comp. Quince seeds $\frac{3}{2}$ ij, water $\frac{3}{2}$ xx.

Dose. f $\frac{3}{4}$ ij-iv, ad libitum.

Incomp. Alcohol, acids, metallic solutions.

FICUS. (F. carica. The dried fruit. Figs.)

Form. Unaltered; decoction, cataplasm.

GLYCYRRHIZA. (G. glabra. Common Liquorice. The root.)

Form and Dose. Decoction (rad. cont. $\frac{3}{4}$ iss, water $\frac{3}{4}$ xx), ad libitum.

1. EXTRACTUM GLYCYRRHIZÆ.

Form and Dose. In substance and mixture. Ad libitum. (Gr. x. covers the taste of a black dose.)

HORDEUM. (H. distichon. Barley. Decorticated seeds.)

1. DECOCTUM HORDEI. (Barley water.)

Comp. Barley $\frac{3}{4}$ ij, water Oivss, boil to Oij.

Dose. Ad libitum.

2. DECOCTUM HORDEI COMPOSITUM.

Comp. Decoct. hord. f $\frac{3}{4}$ xl, fici concisi $\frac{3}{4}$ iiss, glycyrr. rad. 3 v, uvar. pass. $\frac{3}{4}$ iiss, aquæ f $\frac{3}{4}$ xx, boil to f $\frac{3}{4}$ xx.

Dose. Ad libitum.

LINUM. (L. Usitatissimum. Common Linseed.)

The seeds.)

1. OLEUM LINI. (Linseed oil.)

Form. Enema, liniment.

2. INFUSUM LINI COMPOSITUM. (Linseed tea.)

Comp. Lin. sem. $\frac{3}{4}$ ss, glycyrr. rad. 3 ij, water Oj.

Dose. f $\frac{3}{4}$ ij-iv, or ad libitum.

Incomp. Alcohol. Acetate of lead.

3. CATALPLASMI LINI. (See PART II.)**MALVA.** (M. Silvestris. Common Mallow.)**1. DECOCTUM MALVÆ COMPOSITUM.***Comp.* Mallow dried $\frac{2}{3}$ j, chamomile dried $\frac{2}{3}$ ss, water f $\frac{2}{3}$ xx.*Form.* Enema, fomentation.**MEL.** (Honey.)**1. OXYMEL.** (Mel. acetatum. Simple oxymel.)*Comp.* Honey $\frac{2}{3}$ xij, acetic acid f $\frac{2}{3}$ iiij.*Dose.* f $\frac{2}{3}$ j - $\frac{2}{3}$ j.**OLIVÆ OLEUM.** (Olive Oil.)*Form.* Enema, liniment.**TRAGACANTHA.** (Concrete juice of Astragalus Tragacantha.)*Form and Dose.* Powder or mixture, ad libitum.**1. PULVIS TRAGACANTHÆ COMPOSITUS.***Comp.* 1 tragacanth, 1 starch, 1 pulv. acaciæ, 2 sugar.*Dose and Form.* $\frac{2}{3}$ j - 3 ij, in powder and solution.**SARSAPARILLA.** (See ALTERATIVES.)**TAMARINDUS.** (See REFRIGERANTS.)*Diuretics.***CANTHARIS?** (Cantharides. Spanish fly.)
(See STIMULANTS.)*Form and Dose.* Pill, gr. ss-j-ij. (Rarely used.)*Use.* In dropsy when the system is in an atonic condition.

1. ACETUM CANTHARIDIS. (See PART II.)

2. TINCTURA CANTHARIDIS.

Comp. Canth. $\frac{3}{2}$ i, dil. alc. Oij.*Dose.* ℥x- $\frac{3}{2}$ j, in demulcent liquids.

3. CERATUM CANTHARIDIS. (See PART II.)

4. EMPLASTRUM CANTHARIDIS. (See PART II.)

CHIMAPHILA? (*Pyrola Umbellata.* Leaves of the Winter Green.)*Use.* In dropsy attended with disordered digestion and general debility.

1. DECOCTUM CHIMAPHILÆ.

Comp. Chimaphila $\frac{3}{2}$ j, water Oiss, (boil to Oj).*Dose.* $\frac{3}{2}$ j-iij.*Incomp.* Sulphas ferri.**COLCHICUM?** (*C. autumnale.* Cormus and seeds.) (See PURGATIVES.)**DIGITALIS.** (*D. purpurea.* The leaves and seeds.)*Dose.* Gr. ss-iss, in powder or pill.*Use.* In dropsies when the action of the heart requires to be controlled.

1. INFUSUM DIGITALIS.

Comp. Digitalis $\frac{3}{2}$ j, water Oss, tinct. of cinnamon $\frac{3}{2}$ j.*Dose.* $\frac{3}{2}$ ss-j.*Incomp.* Sulphate of iron, acetate of lead.

2. TINCTURA DIGITALIS.

Comp. Digitalis $\frac{3}{2}$ iv, dil. alc. Oij.*Dose.* ℥x- $\frac{3}{2}$ ss-j.

3. EXTRACTUM DIGITALIS. (Watery extract from the fresh leaves.)

Dose. Gr. ss-ijj.

[*Note.*—Pilulæ digitalis et scillæ.—ED.

Comp. Digitalis and squill àà gr. j, conf. arom. gr. ij, confect. rosæ q. s. M. ft. pil.]

DIOSMA. (See DIAPHORETICS.)

DULCAMARA ? (Solanum D. Woody Night-shade. Bitter sweet. The stem.) See DIAPHORETICS. NARCOTICS.)

Use. Mostly in cutaneous eruptions.

1. DECOCTUM DULCAMARÆ.

Comp. Dulcamara ʒ j, water Oiss, boil to Oj.

Dose. f ʒ j-iv.

JUNIPERUS. (Common Juniper. Tops and berries.)

Form and Dose. Berries (triturated with sugar) ʒ j-ij. Infus. (J. ʒ j, water ʒ xv) ʒ ij-iv.

Use. In dropsy.

1. OLEUM JUNIPERI.

Dose. ʒ ij-vj, in pill or emulsion.

2. SPIRITUS JUNIPERI COMPOSITUS.

Comp. Juniper berries ʒ j, caraway and fennel àà ʒ iss, dil. alc. 1 gall., water Oij. Distil to 1 gall.

Dose. f ʒ ij- ʒ ss, (an adjunct).

PAREIRA ? (Cissampelos P. Pareira brava. The root.)

Dose. ʒ ss-j of the powder.

Use. Chronic diseases of the urinary passages.

1. INFUSUM PAREIRÆ.

Comp. Pareira 3vj, water 3 xx.

Dose. 3 j-iij, (used with the extract).

2. EXTRACTUM PAREIRÆ. (Watery extract.)

Dose. Gr. x-3 ss.

POTASSA. (ANTACIDS. ALKALINES.)

1. ACETAS POTASSÆ.

Dose. 3j-3 iss, in diluents.

Incomp. Most acids and neutral salts.

2. CARBONAS POTASSÆ. (Olim. POT. SUBCARB.)

Dose. Gr. x-3 ss.

[NOTE.—3j is saturated by acid. cit. gr. xvij,
acid. tart. gr. xvij, succ. lim. 3 ss.]

3. BICARBONAS POTASSÆ. (Olim. POT. CARE.)
(Milder than 2.)

Dose. Gr. x-3 ss.

[NOTE.—3j is saturated by acid. cit. gr. xiv,
acid. tart. gr. xv, succ. lim. 3 iiiss.]

4. LIQUOR POTASSÆ CARBONATIS.

Comp. Carb. pot. 1bj, water dist. f 3 xij.

Dose. ℥xx-3 j.

Incomp. Acids, acidulous salts, &c.

5. NITRAS POTASSÆ. Nitre. (See REFRIGERANTS.)

Dose. Gr. v-3 ss.

Incomp. Sulphuric acid and its salts.

SCILLA. (S. Maritima. Squills. Recent bulb.)

(See EMETICS. EXPECTORANTS.)

Dose. Gr. v-xv.

Use. Acts best in absence of inflammatory excitement.

1. PULVIS SCILLÆ. (Not officinal, but in use.)

The dried bulb powdered.

Dose. Gr. j-v, in pill combined.

2. PILULÆ SCILLÆ COMPOSITÆ.

Comp. Squills 3j, ginger and gum ammoniac,
each 3ij, soap 3iiij, syr. q. s. Div. in 120 pills.

Dose. Gr. v-x.

3. TINCTURA SCILLÆ.

Comp. Squill $\frac{2}{3}$ iv, proof spir. Oij.

Dose. $\frac{1}{3}$ x-3 ss.

4. ACETUM SCILLÆ.

Comp. Squill $\frac{2}{3}$ iv, distilled vinegar Oij, alc. f $\frac{2}{3}$ j.

Dose. f $\frac{2}{3}$ ss-ij in aromatic water.

Incomp. Alkalies, earths, and carbonates.

5. OXYMEL SCILLÆ.

Comp. Honey $\frac{1}{2}$ iiij, vinegar of squill Oij.

Dose. 3j-ij.

SCOPARIUS. (Scytisus S. De Candolle.

Broom. The fresh tops.)

1. INFUSUM SCOPARII.

Comp. Scoparius $\frac{2}{3}$ j, water $\frac{2}{3}$ xx. [Decoction
preferable. Scop. rec. f $\frac{2}{3}$ ss, water f $\frac{2}{3}$ xx, boil
to $\frac{2}{3}$ x.]

Dose. f $\frac{2}{3}$ j-ij

2. DECOCTUM SCOPARII COMPOSITUM.

Comp. Fresh broom-tops, juniper berries, dande-
lion-root, $\ddot{\text{aa}}$ $\frac{2}{3}$ ss, water f $\frac{2}{3}$ xxx. Boil to $\frac{2}{3}$ xx.

Dose. f $\frac{2}{3}$ j-ij.

SENEGA? (See DIAPHORETICS. STIMULANTS.)

Dose. Of the powder (rarely used) gr. x-xx.

1. DECOCTUM SENEGÆ.

Comp. Senega root $\frac{2}{3}$ j, water Oiss, boil to Oj.

Dose. f $\frac{2}{3}$ j-ij.

[**EXTRACTUM SENEGRÆ.** (Not officinal.)

Dose. Gr. v-x.]

SPIRITUS ÆTHERIS NITRICI. (Sp. of Nit. Ether. Sweet spirits of nitre.)

Dose. 3 ss-ij.

Incomp. Sulphas ferri. Tinct. guaiaci.

Use. In febrile complaints with deficient urinary secretion.

TABACUM. (*Nicotiana T.* Tobacco. The leaves.) (**NARCOTICS.** **PURGATIVES.**)

1. **ENEMA TABACI.** **INFUSUM TABACI.** (See PART II.)

TARAXACUM? (Root of the dandelion.) (See **TONICS.** **ALTERATIVES.** **PURGATIVES.**)

Use. Said to have a specific action on the liver.

1. **EXTRACTUM TARAXACI.** (Watery extract of fresh root.) Should be brown and bitter.

Dose. Gr. x-3 ss-j.

[*NOTE.—Decoctum Taraxaci.—Edinb.* Tar. 3 iiiss, water 3 xx (to 3 x.) *Dose.* 3 j-ij. U. S. Tarax. rad. 3 ij, water Oij. *Dose.* f 3 ij.]

TEREBINTHINA. (Turpentine. Oils and resins from various species of the genus *Pinus*.)

(See **PURGATIVES.** **ANTHELMINTICS.**)

1. **TEREBINTHINA VULGARIS.** (Common turpentine.)

Dose and Form. 3j-3j. (Not often internally.) In pill, emulsion, electuary.

2. **TEREBINTHINA CANADENSIS.** (Liquid turpentine. Canada balsam.)

3. TEREBINTHINA CHIA. (Chian Turpentine. Liquid resin. From the Pistacia Terebinthus.)

4. OLEUM TEREBINTHINÆ.

Dose and Form. ℥vj-3j. (Floating on aromatic water with tinct. capsici, in emulsion with mucilage or white of egg, or as a *Linctus* with honey or aromatic syrup.)

UVA URSI? (See ASTRINGENTS, TONICS?)

1. DECOCTUM UVÆ URSSI.

Comp. Uva ursi ʒj, water f ʒ xx, boil to Oj.

Dose. f ʒ j-iij.

2. EXTRACTUM UVÆ URSSI.

Dose. Gr. v-xv.

Use. In chronic urinary disorders.

Emetics.

AMMONIÆ SESQUI-CARBONAS. (Ammonia Subcarbonas.)

Dose and Form. ʒ ss in any aqueous medium.
(Rarely used.)

ANTHEMIS? (See AROMATIC BITTERS.)

1. INFUSUM ANTHEMIDIS. Chamomile tea.

Comp. Cham. ʒ ss, water Oj (ten minutes).

Dose. ʒ iv subinde (warm).

Use. As an adjuvant.

ANTIMONIUM.

1. ANTIMONII POTASSIO-TARTRAS. (Antimonium Tartarisatum.)

Dose. Gr. j-iv in solution.

Use. A certain, powerful, and permanent emetic.

Used where it is required not only to evacuate the stomach, but to produce an effect on surrounding organs.

2. VINUM ANTIMONII POTASSIO-TARTRATIS. (Vinum Antimonii.)

Comp. 3ij of ant. pot. tart. in 3 xx of sherry.

Dose. 3 ss-iss subinde (gr. j in 3 ss).

Incomp. Bitter and astringent infusions. Alkalies and their carbonates.

Use. For infants.

ASARUM. (Asarum Europaeum. Asarabacca.)

Dose. Root or leaves in powder, 3 ss-j. (Rarely used.)

CUPRI SULPHAS. (Blue vitriol.)

Dose. Gr. v-xv, in solution.

EUPHORBIUM. (Gum resin. Spurge.) (Rarely used.)

IPECACUANHA. (See DIAPHORETICS.)

1. PULVIS IPECACUANHÆ.

Dose. Gr. xv-3 ss in aqueous fluids.

Use. Mild and efficient.

2. VINUM IPECACUANHÆ.

Comp. 3 ij of ipec. in Oij of sherry wine.

Dose. f 3 ss-j every ten minutes. (For children.)

LOBELIA. (Lobelia inflata. Indian tobacco.)

Dose. In powder, gr. xv-3j; of tincture (lobelia 3 iv, dil. alc. Oij), 3 j-ij. (Rarely used.)

Use. In spasmodic asthma.

SCILLA. (Squills.) (See DIURETICS, NARCOTICS, EXPECTORANTS.)

Dose. Gr. v-xv. (Rarely used.)

SINAPIS. (S. nigra. Mustard.) (EXPECTORANTS.)

Dose. (Of the powder) $\frac{3}{j}$ - $\frac{2}{j}$ ss, in $\frac{2}{3}$ iv-vj of warm water.

Use. In narcotic poisoning.

TABACUM. (Tobacco.) (NARCOTICS.)

Dose. Gr. v-x. (Rarely used.)

Use. In reduction of dislocations.

VERATRUM. (White Hellebore.) (See CATHARTICS.)

Form and Dose. Powder, gr. j-vj.

1. VINUM VERATRI.

Comp. White Hellebore $\frac{2}{3}$ iv, sherry Oj.

Dose. \mathfrak{M} x-xxx.

ZINCI SULPHAS. (White vitriol.)

Dose. Gr. xv- $\frac{3}{j}$ ss, in solution.

Use. A prompt emetic in cases of poisoning.

Emmenagogues.

(MENSTRUAE ET PARTUM ACCELERANTIA.)

ERGOTA. (Secale Cornutum. Spurred Rye.) (NARCOTICS?)

Dose. Powder (Pulvis Ergotæ) gr. v-x-xx.

[a. *Infusum.* Ergot $\frac{3}{j}$, boiling water $\frac{2}{3}$ iv.

Dose. $\frac{2}{3}$ iss.

- b. *Decoctum.* Ergot 3 j, wat. 3 v, boil 10 min.
Dose. 3 ij.
- c. *Tinctura.* Ergot 3 ss, spir. rect. 3 vj.
Dose. 3 j.
- d. *Vinum.* (Amer. Ph.) Ergot 3 j, wine 3 viij, macerate 14 days.]

RUTA. (Ruta Graveolens. Rue. The leaves.)

Dose. Powder 3 j-3 ss; infusion (Rue 3 j, boiling water 3 xx) 3 j-ij.

1. CONFECTIO RUTÆ.

Comp. Rue, caraway, bay berries (Laurus nobilis) aa 3 iss, sagapenum 3 ss, black pepper 3 ij, honey 3 xvij. (See CARMINATIVES.)

Dose. 3 j. (Also as an enema for children.)

SABINA. (Juniperus S. Savin. The tops.)

Dose. Powder gr. v-x. (Infusion. Oil.)

From other Classes.

ALOE.

ASSAFÆTIDA.

CANTHARIS.

CASTOREUM.

FERRUM.

GALBANUM.

HELLEBORUS NIGER.

HYDRARGYRUM.

IODINUM.

SENEGA.

SINAPIS.

Errhines.

ASARUM. (See EMETICS.)

EUPHORBIUM. (See EMETICS.)

HELLEBORUS. (See EMETICS.)

TABACUM. (See NARCOTICS.)

Expectorants.

ALLIUM? A. sativum. Garlic. The bulb.
(STOMACHIC, DIURETIC.)

Dose. 3 j-ij of the fresh bulb.

AMMONIACUM. (See ANTISPASMODICS.)

Use. In deficient expectoration without acute inflammation.

ANTIMONII POTASSIO-TARTRAS. (See EMETICS, DIAPHORETICS.)

Dose. Gr. $\frac{1}{2}$ — $\frac{1}{4}$.

Use. Excites bronchial secretion.

ASSAFAETIDA. (See ANTISPASMODICS.)

BALSAMUM PERUVIANUM. (Balsam from the incised bark of the Myroxylon Peruferum. Liquid Balsam.)

Dose and Form. 3 ss-j on sugar, in pill with powders, mucilage.

Use. In chronic catarrh.

BALSAMUM TOLUTANUM. (Concrete Balsam.)

Dose and Form. Same as B. Peruvianum.

1. TINCTURA BALSAMI TOLUTANI.

Comp. Tolu $\frac{3}{4}$ iij, alcohol Oij.*Dose.* $\frac{3}{4}$ ss-ij. (Water precipitates the resin.)

2. SYRUPUS TOLUTANUS.

Comp. Tinc. tolu $\frac{3}{4}$ j, syrup Ojss.*Dose.* f $\frac{3}{4}$ j-ij. (To flavour.)

BENZOINUM. (See ANTISPASMODICS.)

COPAIBA. (Liquid resin.)

Dose and Form. M x- $\frac{3}{4}$ j, on sugar, in water, in emulsion, in syrup, in pill, in capsules.*Use.* In chronic catarrh.

GALBANUM. (See ANTISPASMODICS.)

Use. Chronic bronchial affections.

IPECACUANHA. (See EMETICS.)

Dose. Gr. ss.

MARRUBIUM ? (M. Vulgare. Horehound.)

(TONICS.)

[Horehound Tea : Marr. $\frac{3}{4}$ j, water $\frac{3}{4}$ xx.]

MASTICHE. (See ANTISPASMODICS.)

MYRRHA. (Balsamodendron M. Gumresinous exudation.) (TONICS.)

Form and Dose. Powder, pill, emulsion, gr. x- $\frac{3}{4}$ ss.

1. TINCTURA MYRRHÆ. (Seldom used internally.)

STYRAX. (S. officinale. The balsam.) (ANTISPASMODICS.)

Dose. Gr. v-x-xx, in pill (purified).

1. PILULÆ STYRACIS COMPOSITÆ. (NARCOTICS. OPIUM.)

Comp. 3 storax, 1 opium, 1 saffron.

Dose. Gr. iss-iiss-x (gr. v contains 1 gr. opium).

SAGAPENUM. (See ANTISPASMODICS.)

SENEGA. (See DIAPHORETICS.)

SCILLA. (See DIURETICS.)

CHLORINUM. (Inhaled.) In phthisis.

IODINUM. (Inhaled.) In phthisis.

MYRRHA. (Smoked.) In asthma.

PIX LIQUIDA. (Smoked.) In asthma.

STRAMONIUM. (Smoked.) In asthma.

TABACUM. (Smoked.) In asthma.

SYR. SCILL. COMP. (Coxe's Hive Syrup, U. S.)

Comp. Scill., seneka ää ʒ iv, ant. tart. grs. xlviij, water Oiv, sugar ℥iijss.

Dose. For a child, ℥x-f ʒ j, repeated in cases of croup every fifteen minutes, till it vomits.

Use. It is emetic, diaphoretic, expectorant, and frequently cathartic.

Narcotics. *Sedatives.* *Soporifics.*

Anodynes.

ÆTHER. (See STIMULANTS..)

ACIDUM HYDROCYANICUM DILUTUM.

Comp. Real hydrocyanic acid and water.

Dose. ℥iij-v-vijj, in aqueous mixtures.

Incomp. Metallic oxides and carbonates.

[NOTE.—The dilute acid of the Lond. P. contains 2 per cent. ; of the Dub. 1·6 per cent. ; and of the Ed. 3·2 per cent. of the real acid. The acid of *Scheele's strength* varies ; it is often twice or thrice as strong as the Lond. dilute acid. It is to be desired that the profession would uniformly prescribe the dilute acid of the Lond. Ph. : it keeps better than Scheele's, and the dose is more manageable.]

Use. In asthma, pertussis, chronic catarrh. As a lotion in cutaneous diseases, for itching.

ACONITUM. (A. Paniculatum. Wolfsbane.
Root and leaves.)

Dose. Of the powder, gr. j—iiij. (Not to be relied on.)

Use. In neuralgia, gout, cancer, rheumatism.

1. EXTRACTUM ACONITI. (The inspissated juice.)

Dose. Gr. ss—j—iiij. (Uncertain.)

[*a. Tinctura aconiti.*

Comp. Aconite root, $\frac{3}{4}$ iv, dil. alc. Oij.

Dose. $\frac{M}{xx}$ ter die. (Most valuable as embrocation in neuralgia.)

b. Extractum Alcoholicum Aconiti. (By evaporating the tincture.)

Dose. Gr. $\frac{1}{6}$ — $\frac{1}{2}$, in pill.

[NOTE.—The extract from the root, rubbed down with twice its bulk of alcohol and strained, is the best preparation.]

c. Unguent. Aconiti. One part extract and two parts lard.]

2. ACONITINA.

Dose. Scarcely admissible *internally*. (See PART II.)

ALCOHOL. (See STIMULANTS.)

BELLADONNA. (*Atropa B.* Deadly Night-shade. The leaves.)

Dose. Of the powder, gr. j-v. (The extract and tincture preferable.)

Use. In neuralgia, &c.

1. **EXTRACTUM BELLADONNAE.** (The inspissated juice.)

Dose. Gr. $\frac{1}{2}$ -iv. (More used externally.)

2. **EMPLASTRUM BELLADONNAE.**

(See PART II.)

[*NOTE.*—Though not officinal, the following is useful :

Tinctura Belladonnæ.

Comp. Leaves of bell. $\frac{3}{4}$ iv, dil. alc. Oij.

Dose. M_{xx}-xxiv : (or saturated tincture, extract. bell. $\frac{3}{4}$ x, proof spir. f $\frac{3}{4}$ xij. *Dose.* M_{ij}-iiij-vj.)

CAMPHORA. (*Laurus C.* A peculiar concrete.)

Dose. Gr. j-xv, in pill, or (better) emulsion.

1. **MISTURA CAMPHORÆ.**

Comp. Camphor 3 ss, rect. spir. M_x, water $\frac{3}{4}$ xx, ($\frac{3}{4}$ j = rr. iss of camphor).

Dose. $\frac{3}{4}$ j-ij, or ad libitum.

Use. In typhoid diseases, rheumatism, dysmenorrhœa, strangury.

2. **TINCTURA CAMPHORÆ.** (Camphorated spirits of wine.)

Comp. Camph. $\frac{3}{4}$ iv, alcohol Oij.

Dose. ℥x-3 ss, (in saccharine or mucilaginous fluid; also externally).

3. **TINCTURA CAMPHORÆ COMPOSITA.** (Paregoric elixir. Tinct. opii camphorata.)

Comp. Opium and benzoic acid, each 3 j, Ol. anis. 3 j, mel. desp. 3 ij, camph. 3 ij, dil. alc. Oij.

Use. Relieve nausea, check diarrhoea. Infantile diseases, to procure sleep.

Dose. 3 ss-3 ij.

4. **LINIMENTUM CAMPHORÆ.** (See PART II.)

5. **LINIMENTUM CAMPHORÆ COMPOSITUM.** (See PART II.)

[**CANNABIS INDICA.** (Indian Hemp.) (Not officinal.)

1. **EXTRACTUM CANNABIS INDICÆ.**

Dose. Gr. ss-ij.]

COLCHICUM. (PURGATIVES. DIURETICS.)

CONIUM. (C. maculatum. The leaves and fruit.)

Dose. Of the powder gr. ij-x.

Use. In cancer, pertussis, scrofulous tumours.

1. **TINCTURA CONII.**

Comp. Hemlock 3 iv, dil. alc. Oij.

Dose. 3 ss-j.

2. **EXTRACTUM CONII.** (The inspissated juice.) (Often inert.)

Dose. Gr. ij-xx.

3. **PILULÆ CONII COMPOSITÆ.**

Comp. 5 ext. conii, 1 ipecacuan.

Dose. Gr. v-x (6 grs.=gr. j ipec).

4. CATAPLASMA CONII. (See PART II.)

DIGITALIS. (DIURETICS.)

DULCAMARA. (DIURETICS, DIAPHORETICS, q. v.)

HELLEBORUS. (CATHARTICS, EMETICS, q. v.)

HYOSCYAMUS. (H. niger. Henbane. The leaves and seeds.)

Dose. Powder (rarely used) gr. iij-x.

Use. In cases where opium is inadmissible from constipating effects.

1. EXTRACTUM HYOSCYAMI. (Inspiss. juice.)

Dose. Gr. v-x- 3j.

2. TINCTURA HYOSCYAMI.

Comp. Henbane 3 iv, dil. alc. Oij.

Dose. 3 ss-ij.

LACTUCARIUM. (Lactuca sativa. Lettuce. The inspissated juice.)

Dose. Gr. ij-vj.

Incomp. Ammonia, alcohol.

Use. Occasionally instead of opium.

LACTUÆ EXTRACTUM. (Watery Extract of the same.)

Dose. Gr. v-xv.

LOBELIA. (See EMETICS, ANTISPASMODICS.)

LUPULUS. (See AROMATIC BITTERS.)

NUX VOMICA. (Strychnos Nux Vomica. Ratsbane. The seeds.)

Dose. Gr. iij-xij.

1. STRYCHNIA. (Six times as strong as the alcoholic extract of nux vomica.)

Dose. Gr. $\frac{1}{16}$ -j.

[*NOTE.*—1. *Tinctura Nucis Vomicæ.* (Dub.)
Comp. Nux vom. 3 ij, rect. spir. 3 viij.

Dose. M v-x.

2. *Extractum Nucis Vomicæ.* (Ed. Dub.)
Spirituous extract.

Dose. Gr. ss-j-ijj.

3. Brucia. (From the nux vomica bark.)

Dose. Gr. ss-v.

OPIUM (and MORPHIA.) (Papaver Somniferum. The concrete juice of the unripe capsules.)

Dose and Form. Pill, powder, lozenge, electuary,
gr. $\frac{1}{4}$ -ss-j-ij-v-x.

Use. A general narcotic.

1. PILULÆ SAPONIS COMPOSITÆ.

Comp. Opium 3 ss, soap 3 ij.

Dose. Gr. iss-iiss-x (gr. v=opii gr. j).

2. PILULÆ STYRACIS COMPOSITÆ.

Comp. 1 opium, 1 saffron, 3 storax.

Dose. Gr. iss-iiss-x (gr. v=opii gr. j).

3. EXTRACTUM OPII PURIFICATUM (Watery extract:
 $\frac{1}{3}$ or $\frac{1}{4}$ th more active than opium.)

Dose. Gr. $\frac{1}{4}$ -iv.

4. TINCTURA OPII. (Laudanum.)

Comp. Opium 3 iiiss, dil. alc. Oij.

Dose. M v-3 j.

5. VINUM OPII. (Sydenham's liquid laudanum.)

Comp. Opium 3 ii, cinnamon 3 j, cloves 3 j,
sherry Oj.

Dose. $\frac{1}{2}$ v-3 j, (about the same strength as tinct. opii, but more agreeable to taste).

6. CONFECTIO OPII.

Comp. Opium 3 ivss, aromatic powder, $\frac{3}{2}$ vj, mel. $\frac{3}{2}$ xiv (36 grs.=opii. gr. j).

Dose. Gr. x-3 j.

Use. In diseases requiring a stimulating narcotic.

7. ENEMA OPII. (See PART II.)

8. LINIMENTUM OPII. (See PART II.)

9. TINCTURA CAMPHORÆ COMPOSITA. (See CAMPHOR.)

Comp. Camphor, opium, benzoic acid, &c.

Dose. f 3 ss- $\frac{3}{2}$ ss (f 3 j=opii gr. ij nearly).

10. PULVIS IPECACUANHÆ COMPOSITUS. (DIAPHORETICS.)

Comp. 1 opium, 1 ipecac., 8 sulph. potassæ.

Dose. Gr. iiiss-v- $\frac{1}{2}$ j (gr. x=opii gr. j).

11. PILULÆ IPECACUANHÆ COMPOSITÆ. (DIAPHORETICS.)

Comp. 3 pulv. ipec. c., 1 squills, 1 ammoniac.

Dose. Gr. v-x (gr. v=gr. $\frac{1}{3}$ opium).

13. PULVIS KINO COMPOSITUS.

(ASTRINGENTS.) (One gr. of opium in $\frac{1}{2}$ j.)

13. PULVIS CRETÆ COMPOSITUS CUM OPIO.

(ANTACIDS.) (One gr. of opium in $\frac{1}{2}$ ij.)

14. ACETAS MORPHIÆ.

Dose. Gr. $\frac{1}{2}$ -ss-j.

Incomp. Mineral acids, alkalies.

15. HYDROCHLORAS MORPHIÆ. (Muriate of morphia.)

Dose. Gr. $\frac{1}{2}$ -ss-j.

[*Morphiæ muriatis solutio.* ED. PH.—Hydr. morphiæ, gr. xvij, spir. rect. $\frac{3}{4}$ j, water $\frac{3}{4}$ iiij.
Dose. $\frac{1}{2}$ ij—xxx.

Trochisci morphiæ. ED. PH.—Hydr. morph. gr. v, tinct. tolu $\frac{3}{4}$ j, sugar $\frac{3}{4}$ vj, = 130 lozenges, each containing gr. 1-26 of morphia.

Morphia itself is little used from its insolubility.

Liquor Opii Sedativus. Battley's sedative.

Comp. Said to be a solution of the watery extract.

Dose. $\frac{1}{2}$ ij—xx.

Solutio Morphiæ Bimeconatis. This valuable preparation was introduced by Mr. Squire, in 1838. It is of the same strength as laudanum. It usually disorders the head less than other preparations of opium, and does not constipate the bowels.

Dose. $\frac{1}{2}$ ij— $\frac{3}{4}$ j,

Acetum Opii. Substitute for black drop.

Comp. Opium $\frac{3}{4}$ viij, distilled vinegar q. s. nutmeg $\frac{3}{4}$ iss, saffron $\frac{3}{4}$ ss, sugar $\frac{3}{4}$ xij.

Dose. $\frac{1}{2}$ ij—xx.]

Use. Does not produce so much nausea and headache as laudanum.

PAPAVER. (P. somniferum. The ripe capsules.)

1. DECOCTUM PAPAVERIS. (See PART II.)

2. EXTRACTUM PAPAVERIS. (Watery extract.)

Dose. Gr. ij— $\frac{3}{4}$ j.

3. SYRUPUS PAPAVERIS.

Comp. A concentrated decoction and sugar.

Dose. (For infants,) $\frac{3}{4}$ ss—j; (adults,) $\frac{3}{4}$ ij— $\frac{3}{4}$ ss.

SABADILLA. (*Helonias officinalis.* The seeds.)
—Used chiefly to yield **VERATRIA**, q. v.

STRAMONIUM. (*Datura S.* Thornapple.
Leaves and seeds.) (**EXPECTORANTS, ANTI-SPASMODICS.**)

Dose. Leaves gr. j-ij, seeds gr. ss-j.

1. **EXTRACTUM STRAMONII.** (Watery extract.)

Dose. Gr. $\frac{1}{4}$ -j.

[*Tinctura Stramonii.* (Ph. Amer.) *Comp.*
Stram. seeds $\frac{3}{4}$ j, proof sp. $\frac{3}{4}$ viij. *Dose.* M $\ddot{\text{x}}$ -xx.]

TABACUM. (See **EMETICS, DIURETICS.**)

TOXICODENDRON. (*Rhus Toxicodendron.*
Sumach. The leaves.)

Dose. Powdered leaves gr. ss-iiij.

VERATRIA. (Alkaloid from Sabadilla.)

Dose. Gr. 1-12-ss in pill. (Rarely used internally.)

Comp. Veratria $\frac{3}{4}$ j, rect. spir. $\frac{3}{4}$ ij. (External.)

Unguentum Veratriæ. (See **PART II.**)

VERATRUM. (*V. album.* White Hellebore.
The root.) (**EMETICS, PURGATIVES.**)

Dose. Of the powder gr. j-ij.

Use. In lepra and other cutaneous affections.

1. **DECOCTUM VERATRI.**
(*External.*) (See **PART II.**)

2. **UNGUENTUM VERATRI.**
(*External.*) (See **PART II.**)

3. **VINUM VERATRI.**

Comp. Hellebore $\frac{3}{4}$ iv, sherry Oj.

Dose. M $\ddot{\text{x}}$ x-xx.

Purgatives.

(Cathartics, Laxatives, Aperients, Eccoprotics,
Drastics, Cholalagogues.)

ALOE. (A. Spicata. The inspissated juice.)

1. EXTRACTUM ALÖES PURIFICATUM.

Dose. Gr. ss-x.

Use. Habitual costiveness, amenorrhœa, ascarides.

[*NOTE.*—An invaluable aperient in small doses and when minutely divided. The “Pilulæ ante cibum” of the Parisian Codex, is an excellent form; viz. Al. opt. gr. xv, Mastiches et Fol. ros. rub. exsic. àā gr. v; Syrup. Absinth. q. s.; Divide in pil. xv. But a dose one half as strong often answers admirably; e. g. Rx. Ext. Al. pur. gr. x; Mastiches et Fol. Ros. rub. exsicc. àā gr. xv; Ol. Absinthii ℥x, M. ft. pil. xx. To be taken *cum prandio.*]

2. PULVIS ALÖES COMPOSITUS.

Comp. 3 aloes, 1 guaiacum, 2 pulvis cinnam.
comp.

Dose. Gr. v-x.

Form. Powder, pill.

3. PILULÆ ALÖES COMPOSITÆ.

Comp. 2 aloes, 1 extract of gentian, ol. carui et
syrup. q. s.

Dose. Gr. j-x.

4. PILULÆ ALÖES CUM MYRRHA. (Rufus' Pills.)

Comp. 2 aloes, $\frac{1}{2}$ saffron, 1 myrrh, syr. q. s.

Dose. Gr. j-x-xv.

[PILULÆ ALÖES ET FERRI. (Ed. Ph.)

Comp. 2 aloes, 3 ferri sulph., 6 pulv. cinnam.
comp., 8 confect. roses.

Dose. Gr. v-x.]

Use. In constipation with debility, suppressed catamenia.

5. PILULÆ CAMBOGLÆ COMPOSITÆ.

Comp. 3 aloes, 2 camboge, 1 ginger, 4 soap.

Dose. Gr. iij-x-xv.

Use. An active purge.

6. PILULÆ SAGAPENI COMPOSITÆ. (Baume de vie.)

Comp. 16 sagapenum, 1 aloes, syrup of ginger q. s.

Dose. Gr. v-x-xv.

7. DECOCTUM ALÖES COMPOSITUM.

Comp. Extract of liquorice 3 vij, carbonate of potass 3 j, aloes 3 jss, myrrh 3 jss, saffron 3 jss, water f $\frac{2}{3}$ xxx (boil down to f $\frac{2}{3}$ xx), compound tinct. of cardam. f $\frac{2}{3}$ vij.

Dose. f $\frac{2}{3}$ ss-ij.

Form. In other fluids, or simple.

Incomp. Acids, salts, &c. (See CARBONATE OF POTASS.)

8. TINCTURA ALÖES.

Comp. Aloes 3 j, extract of liquorice 3 iij, water Oiss, alc. Oss.

Dose. 3 ij- $\frac{2}{3}$ j.

Form. Single, or in other fluids.

9. TINCTURA ALÖES COMPOSITA. (Elix. prop.)

Comp. Aloes 3 iij, saffron $\frac{2}{3}$ j, tincture of myrrh Oij.

Dose. 3 j-ij.

Form. In other liquids, with other purgatives.

Use. Purgative, tonic, emmenagogue.

10. VINUM ALÖES.

Comp. Aloes $\frac{2}{3}$ j, cardam. and ginger each 3 j, sherry Oj.

Dose. $\frac{2}{3}$ ss-ij. (Stomachic 3 j-ij.)

Form. Simple, or in composition.

11. ENEMA ALÖES. (See PART II.)

CAMBOGIA. (Gum resin of Stalagmitis Cambogioides.)

Dose. Gr. j-vj-x.

Form. Pill, powder, solution (with carb. of potass, &c.)—chiefly in composition.

Use. In dropsy with torpid bowels, for tænia.

1. PILULÆ CAMBOGIÆ COMPOSITÆ.

Comp. 2 camboge, 3 aloes, 1 ginger, 4 soap.

Dose. Gr. iij-x-xv.

CASSIA. (Cassia Fistula. The pulp of the pods.)

Dose. $\frac{2}{3}$ ij- $\frac{2}{3}$ j.

Form. Simple, but best in composition with other laxatives. (Not effective.)

1. CONFECTIO CASSIÆ.

Comp. 6 cassia pulp, 2 manna, 1 tamarind pulp, syrup of rose, q. s.

Dose. $\frac{2}{3}$ ij- $\frac{2}{3}$ j. (Not effective—not very eligible.)

2. CONFECTIO SENNAE. (See SENNA.)

COLCHICUM. (The cormus and seeds of Meadow Saffron.)

Use. Said to produce bilious stools.

1. COLCHICI CORMUS.

Dose and Form. Powder, gr. ij-vijj.

2. COLCHICI SEMINA.

Dose and Form. Powder, gr. ij-vijj.

3. TINCTURA COLCHICI.

Comp. Colchicum seeds $\frac{3}{2}$ iv, dil. alc. Oij.

Dose. M_lx-xx- $\frac{3}{2}$ ij.

4. TINCTURA COLCHICI COMPOSITA.

Comp. Colch. seeds $\frac{3}{2}$ j, spir. ammoniæ arom, f $\frac{3}{2}$ xx.

Dose. M_lx-xx- $\frac{3}{2}$ ij.

5. ACETUM COLCHICI.

Comp. Fresh colchicum cormus $\frac{3}{2}$ ij, acet. dist. Oij, alc. f $\frac{3}{2}$ j.

Dose. f $\frac{3}{2}$ ss-ij, in any mild fluid.

Incomp. Alkalies.

6. VINUM COLCHICI.

Comp. Colch. semin $\frac{3}{2}$ iv, sherry Oij.

Dose. M_lx- $\frac{3}{2}$ ss-ij.

7. EXTRACTUM COLCHICI CORMI. (Watery Extract.)

Dose. Gr. j-ijj.

8. EXTRACTUM COLCHICI ACETICUM.

Dose. Gr. j-ijj.

(See ALTERATIVES. EMETICS.)

COLOCYNTHIS. (Cucumis Colocynthis. The dried pulp of the fruit.)

Use. A hydragogue cathartic. Dropsy.

1. COLOCYNTHIDIS PULPA.

Dose. Gr. j-iv-x.

Form. In powder, decoction, or infusion.
manageable, seldom used.)

2. EXTRACTUM COLOCYNTHIDIS. (Watery Extract.)

Dose. Gr. v-x.

Form. Pill. (Keeps badly.)

3. EXTRACTUM COLOCYNTHIDIS COMPOSITUM.

Comp. Coloc. $\frac{3}{2}$ vj, pulv. aloes $\frac{3}{2}$ xij, scammony $\frac{3}{2}$ iv, cardam $\frac{3}{2}$ j, sapo. hisp. $\frac{3}{2}$ iiij; alc. dil. gall. j.

Dose. Gr. ij-v-x.

Form. Pill, single or in combination.

Use. In habitual costiveness.

4. ENEMA COLOCYNTHIDIS. (See PART II.)

ELATERIUM. (A fecula, from the fresh fruit of the Mormordica Elaterium.)

1. EXTRACTUM ELATERII. (Inspissated juice.)

Dose. Gr. $\frac{1}{6}$ -j.

Form. Pill; aqueous solution.

HELLEBORUS. (H. niger. The root.)

Dose. Gr. iij- $\frac{3}{2}$ j. (Rarely used.)

1. TINCTURA HELLEBORI.

Comp. Hellebore $\frac{3}{2}$ iv, dil. alc. Oij.

Form and Dose. Mlxx-3 j. (Rarely used as purgative.)

[**EXTRACTUM HELLEBORI.** (Watery Extract.)]

Dose. Gr. j-iiij.]

HYDRARGYRUM.

1. HYDRARGYRUM CUM CRETĀ.

Comp. 3 mercury, 5 chalk.

Dose. Gr. j-v (alterative), gr. v-xv (laxative).

Form. Powder, pill.

Incomp. Acids and acidulous salts.

Use. For children.

2. HYDRARGYRI CHLORIDUM. Calomel.

Dose. Gr. ss-j (alterative), gr. j-x (purgative).

Form. Pill, powder.

Incomp. Most salts, acids, and alkalies.

Use. In cases where action on liver is desirable.

3. PILULÆ HYDRARGYRI.

Comp. Mercury 1 $\frac{1}{2}$, confect. rosæ 1 $\frac{1}{2}$, liquorice $\frac{1}{2}$.

Dose. Gr. j-iij-v-x.

4. PILULÆ HYDRARGYRI CHLORIDI COMPOSITÆ.

(Plummer's pill.)

Comp. 1 chloride of mercury (calomel), 1 oxy-sulphuret of antimony, 1 treacle, 2 guaiac.

Dose. Gr. iiiss-v-x.

JALAPA. (Ipomæa Jalapa. The root.)

1. PULVIS JALAPÆ.

Dose. Gr. x-3 ss.

Form. Powder, pill, bolus, single or (more generally) combined.

Use. Hydragogue cathartic.

2. PULVIS JALAPÆ COMPOSITUS.

Comp. 1 jalap, 2 bitartrate of potash (cream of tartar).

Dose. 2j-3j.

Form. Powder, mixture.

Incomp. Alkalies and alkaline earths, mineral acids, &c. (See POTASSÆ BITART.)

3. EXTRACTUM JALAPÆ. (Soft and hard.)

Dose. Gr. v-x-xv.

Form. Pill, powder.

4. TINCTURA JALAPÆ.

Comp. Jalap $\frac{3}{2}$ viij, dil. alc. Oij.

Dose. $\frac{3}{2}$ j- $\frac{3}{2}$ ss.

Form. In combination with other purgatives.

MAGNESIA.

1. MAGNESIA. (Calcined magnesia.)

Dose. Gr. x-xx- $\frac{3}{2}$ ss-j.

Form. Powder, electuary, mixture.

Use. In sick headache, acidity of primæ viæ.

2. MAGNESIÆ CARBONAS.

Dose. Gr. xv- $\frac{3}{2}$ ij.

Form. Powder, electuary, mixture.

Incomp. Acids, alkalies, most salts, &c.

Use. In cases requiring an antacid laxative.

3. MAGNESIÆ SULPHAS. (Epsom Salts.)

Dose. $\frac{3}{2}$ ij- $\frac{3}{2}$ iss.

Form. In solution (an oz. of water dissolves an oz. of salt).

Incomp. Muriates, nitrates, acetates, carbonates, &c.

MANNA. (Ornus Europæa. The concrete juice.)

Dose. $\frac{3}{2}$ j-iij- $\frac{3}{2}$ iss.

Form. In solution, in milk, &c.; often with other purgatives.

Use. For children, pregnant women.

POTASSA.

1. POTASSÆ ACETÆ. (See DIURETICS.)

Dose. $\frac{3}{2}$ ij-vj. (Domestic $\frac{3}{2}$ j.)

Form. In solution (an oz. of water dissolves about 3 viss.)

Incomp. Sulphuric, nitric, hydrochloric acids, sulphates, &c.

Use. In dropsies.

2. POTASSÆ SULPHAS.

Dose. Gr. xv-3 j (alt. gr. v-x-xx.)

Form. Powder, solution more rarely, (an oz. of water dissolves 24 grs.).

Incomp. Nitric and hydrocyan. acids, &c.

Use. In visceral obstructions of children.

3. POTASSÆ BISULPHUS. (Potassæ Supersulphas.)

Dose. Gr. x-3 ij.

Form. Powder, solution (soluble in twice its weight of water), combined with rhubarb, &c. (Less used than No. 2.)

Incomp. Alkalies, earths, carbonates, &c.

Use. In constipation, with languid appetite.

4. POTASSÆ TARTRAS.

Dose. 3 j- $\frac{3}{2}$ j.

Form. Solution (soluble when crystallized, in equal weights of water).

Incomp. Most acids and salts.

Use. A cooling purge in febrile diseases.

5. POTASSÆ BITARTRAS. (Cream of Tartar.)

Dose. 3 j-vj.

Form. Solution, electuary, mixture. (Requires 120 parts of water to dissolve it.)

Incomp. Alkalies and alkaline earths, mineral acids, &c.

Use. In small doses, aperient; in large, hydragogue cathartic.

PRUNA. (Prunus domest. The dried fruit.)

Form and Dose. As fruit, dressed or undressed.
Decoction in composition.

RHAMNUS. (Rhamnus Catharticus. Buck-thorn. The berries.)

1. **RHAMNI BACCÆ.** (Baccarum Succus.)

Dose. Recent berries $\frac{1}{2}$ j, dried berries 3j, expressed juice $\frac{3}{4}$ j. (Rarely used.)

2. **SYRUPUS RHAMNI.**

Comp. Buckthorn juice, ginger, pimenta, and sugar.

Dose. 3 iij- $\frac{3}{4}$ j (rarely used); 3j added to purgative draughts and electuaries.

RHEUM. (Rheum Palmatum. Rhubarb. The root.)

Use. Unites astringent with cathartic properties. In dyspepsia with constipation, in diarrhoea if purging indicated, in secondary stage of cholera infantum, in fecal accumulations during typhoid disease.

1. **RHEI PULVIS.**

Dose. Gr. v-xxx.

Form. Powder, pill, electuary, mixture.

2. **INFUSUM RHEI.**

Comp. Rhubarb 3i, water Oss.

Dose. f $\frac{3}{4}$ j-ij. (Stomachic, 3ij-vj.)

Incomp. Stronger acids. Metallic solutions, salts of iron, tart. emet. Inf. of cusparia bark, catechu, &c.

3. **TINCTURA RHEI COMPOSITA.**

Comp. Rhubarb $\frac{3}{4}$ iiss, liquorice 3 vj, ginger 3 iij, saffron 3 iij, proof sp. f $\frac{7}{3}$ xl.

Dose. f 3 ij— $\frac{7}{3}$ iss.

4. EXTRACTUM RHEI.

Dose. Gr. v.—xv.

Form. Pill. Solution in aromatic water.

5. PILULÆ RHEI COMPOSITÆ.

Comp. Rhub. $\frac{3}{4}$ j, aloes 3 vj, myrrh $\frac{3}{4}$ ss, ol. menth. pip. 3 ss, syr. aur. q. s. make 240 pills.

Dose. Gr. v—xv.

[PULVIS RHEI COMPOSITUS. Ed. Ph. (Gregory's Powder.)

Comp. 1 ginger, 2 rhubarb, 6 magnesia.]

RICINI OLEUM. (Ricinus communis. Oil of the seeds. Castor oil.)

Dose. 3 j-iss.

Form. Pure, on some liquid. Emulsion with mucilage, &c.

Use. A mild and speedy cathartic, in fecal accumulations, in colic, diarrhoea, dysentery, enteritis.

ROSA. (R. Centifolia. Damask rose. The petals.)

1. SYRUPUS ROSÆ.

Comp. Sugared decoction.

Dose. 3 ij—vj, (for infants).

SAPO. (See ANTACIDS.)

SCAMMONIUM. (Convolvulus Scammonea. Scammony. The gum resin.)

Dose. Gr. ij—x.

Form. Fine powder triturated with sugar, &c.
In solution.

1. PULVIS SCAMMONII COMPOSITUS.

Comp. 4 scammony, 4 ext. jalap, 1 ginger.

Dose. Gr. v-xv.

Form. Powder, pill, mixture, electuary.

2. CONFECTIO SCAMMONII.

Comp. 2 scammony, 1 cloves, 1 ginger, oil of carui and syrup, q. s.

Dose. Gr. v-x-3j.

3. EXTRACTUM COLOCYNTHIDIS COMPOSITUM. (See COLOCYNTH.)

SENNA. (Cassia lanceolata, and C. obovata. The leaves.)

Dose and Form. Powder 3ss-ij. (Not much used.)

Use. A prompt, efficient, and safe purge.

1. INFUSUM SENNAE COMPOSITUM.

Comp. Senna 3j, coriander 3j, water Oj.

Dose. f 3j-iij. Mostly in composition.

Incomp. Strong acids, metallic salts, lime water.

2. CONFECTIO SENNAE. (Lenitive Electuary.)

Comp. Senna 3viii, figs 1bj, tamarinds 1bss, cassia 1bss, prunes 1bss, coriander 3iv, liquorice root 3iii, sugar 1biiss, water Oiv.

Dose. 3j-iv.

Use. Admirable in habitual costiveness.

3. TINCTURA SENNAE COMPOSITA.

Comp. Senna 3iiiss, caraway 3iiiss, cardamoms 3j, raisins 3v, proof spirit f 3xl.

Dose. 3j-3j.

4. SYRUPUS SENNÆ.

Comp. Senna $\frac{2}{3}$ ij, fennel $\frac{2}{3}$ j, sugar $\frac{2}{3}$ xv, water Oj.

Dose. (For children) f 3 j-ijj.

Use. For children.

SODA.

1. SODÆ PHOSPHAS.

Dose. $\frac{2}{3}$ ss-jss.

Form. Dissolved in gruel or broth.

Use. A mild purgative.

2. SODÆ POTASSIO-TARTRAS. (Soda Tartarisata.)

Rochelle salt.)

Dose. $\frac{2}{3}$ ij- $\frac{2}{3}$ j.

Form. Solution. Single or combined. (Soluble in 5 parts of water.)

Incomp. Most acids and acidulous salts except bitartrate of potass.

Use. A mild, cooling purge.

[*NOTE.* SEIDLITZ POWDERS. *Blue Paper.*

Sod. pot. tart. $\frac{2}{3}$ ij, sodæ bicarb. $\frac{1}{2}$ ij. *White*

Paper. Acid. tart. grs. xxxv.]

3. SODÆ SULPHAS. (Glauber's salt.)

Dose. $\frac{2}{3}$ ij- $\frac{2}{3}$ jss.

Form. Solution. (An oz. of water dissolves $\frac{2}{3}$ iiiss.) Taste disguised by a little lemon juice or bitart. potassæ.

Incomp. Fixed alkalies and their carbonates.

Use. An efficient cathartic.

4. SODÆ ACETAS.

Dose. $\frac{2}{3}$ j- $\frac{2}{3}$ ss.

Incomp. The mineral acids.

SULPHUR. (S. Sublimatum. Flowers of Sulphur.)

Dose. 3 j-ij.

Form. Powder, electuary, suspended in milk, &c.

Use. In haemorrhoids, chronic rheumatism, catarrh, gout.

TAMARINDUS. (Tamarindus Indica. The pulp of the legume.)

Dose. 3 ss-ij, as an adjunct.

Incomp. Salts of soda and potass.

Use. Laxative and refrigerant.

TEREBINTHINÆ OLEUM. (Distilled oil from the resin of Pinus Silvestris.)

1. **TEREBINTHINÆ OLEUM PURIFICATUM.**

Dose. 3 j-ij - 3 iss.

Form. Floating on any fluid. Emulsion.

Use. In ascarides, obstinate constipation, tympanitis.

2. **ENEMA TEREBINTHINÆ.** (See PART II.)

TIGLII OLEUM. (Croton Tiglum. Croton Oil. Expressed from the seeds.)

Dose. ℥ 1-ss-j

Form. Pill, with bread or any extract. Emulsion.

Use. A powerful hydragogue cathartic. Mania, coma, obstinate constipation.

VERATRUM. (White Hellebore.) (See EMETICS.)

Refrigerants.

(Antiphlogistic, Temperants, Febrifuges, Salines.)

ACETUM. (Common Vinegar.)1. **ACETUM DESTILLATUM.** (Distilled vinegar.)*Dose.* f 3 j-ijj, with water and sugar.2. **ACIDUM ACETICUM.** (Seven times stronger than distilled vinegar.)*Dose.* ℥ ij-x, largely diluted.**ACIDUM CITRICUM.***Dose and Form.* Gr. x-⊖j, largely diluted in water, with sugar; or with the alkalies in the form of saline draught.*Incomp.* Acids and alkalies.[*Note.*—The following quantities of alkaline carbonates are required to saturate a scruple of Acid. Cit.: Pot. bicarb. gr. xxix, pot. carb. gr. xxiv, ammon. sesquic. gr. xvij, carb. sodæ gr. xlj, sod. sesquic. gr. xxiv.—*Pereira.*]**ACIDUM HYDROCHLORICUM DILUTUM.***Comp.* Acid. hydrochlor. ʒ j, water f ʒ ijj.*Dose.* ℥ v-xx, largely diluted.**ACIDUM NITRICUM DILUTUM.***Comp.* Acid. nitric. ʒ j, water f ʒ ix.*Dose.* ℥ v-xx, largely diluted.**ACIDUM SULPHURICUM DILUTUM.***Comp.* Acid. sulph. ʒ j, water ʒ xij.*Dose.* ℥ v-xx, largely diluted.*Incomp.* Alkaline carbonates, metallic oxides.**ACIDUM TARTARICUM.**

Dose. 3 ss-3 j, dissolved in much water, also to saturate the alkalies for saline draughts.

Incomp. Alkalies and their carbonates.

ACETOSELLA. (Oxalis A. Wood sorrel.)

Use. Infused in water, or boiled in milk to form whey.

AMMONIA.

1. LIQUOR AMMONIÆ ACETATIS.

Comp. Am. sesquic. 3 ix (*vel* q. s.) acet. dest. f 3 xx. (A saturated solution.)

Dose. f 3 iij-vj.

Incomp. Acids and alkalies.

[AMMONIÆ CITRAS. Amm. sesqui-carb. 3 j. Acid. citric. gr. xxiv, *vel* succ. limon. 3 vj, *vel* acid. tart. gr. xxvj, aquæ et sacch. q. s. (Saline draught.)]

AURANTIUM. (Citrus A. Orange. The fruit.)

LIMONES. (Citrus Limonum. The fruit.)

1. SUCCUS LIMONUM. (Lemon juice.) Diluted as lemonade.

[*Artificial Lemon juice.* Acid. citric. 3 viiss, aquæ destill. 3 xvj, ol. limon. Miv-x.]

2. SYRUPUS LIMONUM.

Comp. Lemon juice 3 x, sugar 3 xv.

Dose. 3 j-iij, added to aqueous fluids.

MORA. (Morus nigra. The Mulberry. The fruit.)

1. SYRUPUS MORI.

Comp. Mulberry juice 3 x, sugar 3 xv.

Dose. 3 j-ij, added to aqueous fluids.

POTASSA.

1. ACETAS POTASSÆ?

Dose. 3j-3i, in diluent mixtures.

2. POTASSÆ BITARTRAS. (Cream of tartar.)

Dose. 3 j-ij in water largely diluted, ad libitum.

(*Imperial*: cream of tartar 3j, boiling water 3 xx, sugar, lemon-peel q. s. *Cream of tartar whey*: 3ij to 3 xx of boiling milk.)

3. POTASSÆ NITRAS. (Nitre.)

Dose. Gr. ij-3j.

Incomp. Sulph. acid, alkaline sulphates.

4. POTASSÆ CARBONAS.

5. POTASSÆ BICARBONAS.

Saline draughts.

[*(a)* Pot. carb. 3j, acid. cit. gr. xvij, *vel* acid. tart. gr. xvij, *vel* succ. lim. 3 ss.

(b) Pot. bicarb. 3j, acid. cit. gr. xiv, *vel* acid. tart. gr. xv, *vel* succ. lim. 3 iiiiss.]

PRUNA. (Prunes. The dried fruit.)

(EMOLLIENTS. PURGATIVES.)

ROSA CANINA. (Dog-rose. The fruit—*Hips*.)

1. CONFECTIO RÖSÆ CANINÆ. (Conserve of Hips.)

Comp. 3 hip-pulp, 5 sugar.

Dose and Form. A vehicle for pills, linctus, or electuary, ad libitum.

RUMEX. Rumex acetosa. Sorrel. The leaves.)

Use. Infused in water, or to form whey.

SODA. (Citrate and tartrate.)

1. SODÆ CARBONAS. (Olim SUBCARBONAS.)
2. SODÆ SESQUICARBONAS. (Olim CARBONAS.)

Saline draughts.

- [*(a)* Sodæ carb. 3j, acid. citric. gr. x ; *vel* acid. tartar. gr. xj ; *vel* succ. limon. 3 iiss.
(b) Sodæ sesquicarb. 3j, acid. citric. gr. xvij ; *vel* acid. tartar. gr. xvij ; *vel* succ. limon 3 ss.]

[PULVERES EFFERVESCENTES. (Ed. Ph.)

Blue Paper. Bicarb. sodæ, gr. xxxij ; *vel* bicarb. pot. gr. xl.

White Paper Acid. tart. gr. xxx.]

TAMARINDUS. (T. Indica. The pulp.)

UVA. (Raisins stoned. The dried fruit.)

Sialogogues. Masticatories.

(See STIMULANTS, ALTERATIVES, NARCOTICS.)

ARMORACIA.

PIPER.

CARYOPHYLLUS.

PYRETHRUM.

HYDRARGYRUM.

TABACUM.

MEZEREUM.

ZINZIBER.

Stimulants.

ÆTHER SULPHURICUS.

Dose. ℥xx-3j.

1. SPIRITUS ÆTHERIS SULPHURICI COMPOSITUS.
(Hoffman's anodyne.)

Comp. Æth. sulph. Oss, spir. rect. Oj, ætherial oil $\frac{3}{4}$ ij.

Dose. $\frac{3}{4}$ ss-j.

Use. To quiet nervous irritation. Added usefully to laudanum to prevent nausea.

ALCOHOL. Sp. grav. .835.

Dose. $\frac{3}{4}$ ss-j.

NOTE.—The words “proof spirit” occurring in the text are to be understood as of sp. grav. 0.920. Such formulæ are unchanged from the London Dispensatory.

2. SPIRITUS TENUIOR. (Dil. alc.) Sp. grav. .935.

Comp. Alcohol and aq. dist., each 1 pint.

Dose. $\frac{3}{4}$ ij- $\frac{3}{4}$ j.

AMMONIA.

1. LIQUOR AMMONIÆ FORTIOR. Sp. grav. .882.
(1 ammonia, 3 water—nearly.)

Dose. $\frac{3}{4}$ v-xv, with water or milk. (Rarely used internally.)

2. LIQUOR AMMONIÆ. (1 ammonia, 9 water.) Sp. grav. .960.

Dose. $\frac{3}{4}$ x-xx.

3. AMMONIÆ SESQUICARBONAS. (Olim A. SUB-CARBONAS.)

Dose and Form. Gr. ij-x, in pill or liquid.

Incomp. Acids and alkalies.

4. LIQUOR AMMONIÆ SESQUICARBONATIS.

Comp. Sesquic. of ammonia $\frac{3}{4}$ j, water $\frac{3}{4}$ v.

Dose. $\frac{3}{4}$ ss-j.

5. SPIRITUS AMMONIÆ.

Comp. Ammonia, carbonic acid, spirit.

Dose. 3 ss-j.

6. SPIRITUS AMMONIÆ AROMATICUS.

Comp. The same, with cinnamon, cloves, and lemon peel.

Dose. 3 ss-j.

Use. In hysteria.

7. SPIRITUS AMMONIÆ FÆTIDUS.

Comp. The same, with assafætida in place of cinnamon. &c.

Dose. 3 ss-j.

8. LINIMENTUM AMMONIÆ. (See PART II.)**9. LINIMENTUM AMMONIÆ SESQUICARBONATIS. (See PART II.)****10. TINCTURA AMMONIÆ COMPOSITA. (Eau de Luce.) (ANTISPASMODICS.)**

Comp. Mastic 3 ij, rect. spir. 3 ix, oil of lavender ℥ xiv, oil of amber ℥ iv, liquor ammon. fort. 3 xx.

Dose. ℥ v-xxx, in water.

Incomp. Acids and earthy salts.

ARMORACIA. (Cochlearea A. Horse-radish. The root.)

Dose. 3 ss-ij, scraped.

Use. In scorbutic affections.

1. SPIRITUS ARMORACIÆ COMPOSITUS.

Comp. Horse-radish, orange peel, nutmegs, proof spirit and water.

Dose. 3 j-ij.

2. INFUSUM ARMORACIÆ COMPOSITUM.

Comp. Horse-radish, mustard, each $\frac{3}{2}$ j, water Oj.
Dose. $\frac{3}{2}$ j-ij.

CAJAPUTI. (*Melaleuca minor.* The oil.) (ANTISPASMODICS.)

Form and Dose. The oil, $\frac{1}{10}$ ij-xx on sugar or in emulsion.

Use. In spasmodic affections of stomach and bowels unconnected with inflammation.

CARDAMINE. (*C. Prætensis.* The cuckoo-flower.)

Dose. $\frac{3}{2}$ j-ij of the dried flowers. (Rarely used.)

CREOSOTON. (Oxyhydrocarburet from pyroxylic oil.)

Dose and Form. $\frac{1}{10}$ j-yj in spirit and water, camphor mixture, or mucilage.

Incomp. Mineral acids.

[*MISTURA CREOSOTI.* (Edin. Ph.)]

Comp. Creosot. et acid. acet. $\frac{1}{2}$ ij xvij, spir. junip. comp. et syrup. $\frac{1}{2}$ ss $\frac{3}{2}$ j, water $\frac{3}{2}$ xiv.

Dose. $\frac{3}{2}$ ss-ij.]

Use. Internally, as an expectorant. Externally to eruptions.

SERPENTARIA. (*Aristolochia S.* Virginian Snakeroot. The root.) (AROMATICS.)

Dose. Powder gr. x- $\frac{3}{2}$ ss.

1 INFUSUM SERPENTARII.

Comp. Serp. $\frac{3}{2}$ ss, water Oj.

Dose. $\frac{3}{2}$ j-ij.

Use. In low forms of fever.

2. TINCTURA SERPENTARII.

Comp. Serp. $\frac{3}{2}$ iii, dil. alc. Oij.

Dose. $\frac{3}{2}$ j-ij.

SPIRITUS VINI GALLICI. (French Brandy.)

Dose. $\frac{3}{2}$ ij- $\frac{3}{2}$ j.

1. MISTURA SPIRITUS VINI GALLICI.

Comp. Ol. cinnamon $\frac{1}{2}$ ij, 2 yolks of eggs, sugar $\frac{3}{2}$ ss, French brandy and cinnamon water $\frac{1}{2}$ $\frac{3}{2}$ iv.

Dose. $\frac{3}{2}$ ij-x. (Egg Flip.)

VINUM XERICUM. (Sherry Wine.)

Dose. $\frac{3}{2}$ ij- $\frac{3}{2}$ i-ij. Ad libitum.

From the Class of AROMATIC BITTERS, q. v.

Absinthium.

Acorus.

Anthemis.

1. Infusum Anthemidis.
2. Oleum Anthemidis.

Aurantii Cortex.

1. Infusum Aurantii compositum.
2. Tinctura Aurantii.
3. Confectio Aurantii.
4. Syrupus Aurantii.
5. Aquæ Florum Aurantii.
6. Oleum Aurantii.

Limonum Cortex.

1. Oleum Limonum.

Contrajervae.

Cusparia.

1. Infusum Cuspariae.

Lupulus.

1. Infusum Lupuli.
2. Tinctura Lupuli.
3. Extractum Lupuli.

*Canella.**Capsicum.*

1. Tinctura Capsici.

Cardamomum.

1. Tinctura Cardamomi.
2. Tinctura Cardamomi composita.

Caryophyllus.

1. Oleum Caryophylli.

Cinnamomum.

1. Pulvis Cinnamomi compositus.
2. Aqua Cinnamomi.
3. Tinctura Cinnamomi.
4. Tinctura Cinnamomi composita.
5. Oleum Cinnamomi.
6. Spiritus Cinnamomi.

Cubeba.

1. Tinctura Cubebæ.

Myristica.

1. Oleum Myristicæ.
2. Spiritus Myristicæ.

Pimenta.

1. Oleum Pimentæ.
2. Spiritus Pimentæ.
3. Aqua Pimentæ.

Piper Longum.

Piper Nigrum.

1. Confectio Piperis Nigri.

Inula.

Sinapis.

Zinziber.

1. Tinctura Zinziberis.
2. Syrupus Zinziberis.
3. Extractum Zinziberis (not officinal).

From NARCOTICS, q. v.

Camphora.

1. Mistura Camphoræ.
 2. Tinctura Camphoræ.
 3. Tinctura Camphoræ composita.
- (Local.) Linimentum Camphoræ.
Linimentum C. compositum.

Dulcamara?

Helleborus?

Nux Vomica. Strychnia?

Toxicodendron?

Veratrum?

From ANTISPASMODICS, q. v.

Castoreum.

1. Tinctura Castorei.

Galbanum.

1. Pilulæ Galbani compositæ,

Moschus.

- 1 Mistura Moschi.

Olibanum.

Opopanax.

Valeriana.

1. Tinctura Valerianæ.
2. Tinctura Valerianæ Ammoniata.

Petroleum.

Succinum.

1. Oleum Succini.

From CARMINATIVES, q. v.

Anethum.

1. Aqua Anethi.

Anisum.

1. Oleum Anisi.
2. Spiritus Anisi.

Carui.

1. Oleum Carui.
2. Spiritus Carui.
3. Aqua Carui.

Coriandrum.

Cyminum.

Lavandula.

1. Oleum Lavandulæ.
2. Spiritus Lavandulæ.
3. Tinctura Lavendulæ composita.

Mentha Piperita.

1. Oleum Menthæ Piperitæ.
2. Spiritus Menthæ Piperitæ.
3. Aqua Menthæ Piperitæ.

Pulegium.

1. Oleum Pulegii.

2. Spiritus Pulegii.
3. Aqua Pulegii.

Mentha Viridis.

1. Oleum Menthæ Viridis.
2. Spiritus Menthæ Viridis.
3. Aqua Menthæ Viridis.

Origanum.

1. Oleum Origani.

Rosmarinus.

1. Oleum Rosmarini.
2. Spiritus Rosmarini.

From EXPECTORANTS, q. v.

Balsamum Peruvianum.

Balsamum Tolutanum.

1. Tinctura Balsami Tolutani.
2. Syrupus Tolutanus.

Myrrha.

1. Tinctura Myrrhæ.

Copaiba.

From EMMENAGOGUES, q. v.

Ergota.

Ruta.

1. Confectio Rutæ.

Sabina.

From DIURETICS, q. v.

Cantharis.

Chimaphila ?

1. Decoctum Chimaphilæ.

Terebinthina Vulgaris.

Terebinthina Canadensis.

Oleum Terebinthinæ.

Spiritus Ætheris Nitrici.

Tonics.

MINERAL TONICS.

(Acids, Earths, Metals.)

ACIDUM ARSENICUM. (White Arsenic.)

Dose. Gr. $\frac{1}{16}$ — $\frac{1}{4}$ in pills with bread, (after a meal.)

Incomp. Acids, earths, bitter infusions.

1. **LIQUOR POTASSÆ ARSENITIS.** (Liquor Arsenicalis. Fowler's Solution.)

Comp. Arsenious acid, carb. of pot. $\ddot{\text{a}}$ $\ddot{\text{a}}$ grs. lxiv, compound tinct. lavender $\frac{3}{2}$ ss, water q. s.

Dose. \mathfrak{M} iij—xx.

Use. In intermitte its, leprosy, cephalalgia.

ACIDUM HYDROCHLORICUM. (A. Muraticum. Liquid hydrochloric acid.)

Dose. \mathfrak{M} v—x in aqueous fluids.

Incomp. Alkalies, earths, &c.

1. **ACIDUM HYDROCHLORICUM DILUTUM.** (Dilute Muriatic Acid.)

Comp. Acid 1 part, water 3 parts.

Dose. \mathfrak{M} x— $\frac{3}{2}$ ss, in aqueous fluids.

(See REFRIGERANTS.)

ACIDUM NITRICUM. (Strong or fuming Nitric Acid.)

Dose. \mathfrak{M} j—x. (Rarely used.)

1. ACIDUM NITRICUM DILUTUM.

Comp. Acid 1 part, water 9 parts.*Dose.* ℥ x-3ss.*Incomp.* Acids, earths, sulph. of iron, &c.

[NOTE. *The Nitro-muriatic bath.* Acid. hydrochlor. 2 parts, acid. nit. 1 part, water 3 parts.— $\frac{3}{4}$ ij of this dilute acid to a gallon of water. (See REFRIGERANTS.)]

ACIDUM SULPHURICUM. (Aqueous Sulphuric Acid. Oil of Vitriol.)

1. ACIDUM SULPHURICUM DILUTUM.

Comp. Acid 1 part, water 13.*Dose.* ℥ x-3ss, in aqueous fluids. As gargle 3 i-ij in $\frac{3}{4}$ viiij.*Incomp.* Earthy oxides, alkaline carbonates.*Use.* For colliquative sweats, hemorrhages.ARGENTI NITRAS. (Argentum Nitratum.)
Lunar Caustic.)*Dose.* Gr. $\frac{1}{6}$ -iiij in pill, with bread crumb or vegetable powder.

1. LIQUOR ARGENTI NITRATIS.

Comp. Argent. nit. 3 j, water $\frac{3}{4}$ j. (A test.)*Use.* Epilepsy, chorea, angina pectoris.

BARII CHLORIDUM. (Muriate of Baryta.)

Used chiefly in the form of—

1. LIQUOR BARII CHLORIDA.

Comp. Chloride of B. $\frac{3}{4}$ i, water $\frac{3}{4}$ iiij.*Dose.* ℥ x-3ss, in distilled water.*Incomp.* Undistilled water, earths, and alkaline salts and sulphates.*Use.* In scrofula and cancer.

BISMUTHI TRISNITRAS?

Dose and Form. Gr. v- $\frac{1}{2}$ j, in pill or powder.

Use. In chronic diseases of the stomach.

CALCII CHLORIDUM. (Muriate of Lime.)

Used in solution as—

1. LIQUOR CALCII CHLORIDI.

Comp. Chloride of lime 1 part, water 2·5 parts.

Dose. ℥ xx-3iss. (See ALTERATIVES.)

Use. In scrofula, tabes mesenterica.

CUPRUM?**1. CUPRI SULPHAS.**

Dose. Gr. $\frac{1}{4}$ -iss, in pill.

2. CUPRI AMMONIO SULPHAS. (C. Ammoniatum.)

Ammoniate of copper.)

Dose. Gr. ss-v, in pill.

3. LIQUOR CUPRI AMMONIO SULPHATIS.

Comp. Ammonio-sulph. of copper 3j, water f $\frac{2}{3}$ xv.

Dose. 3j-ij. Externally as lotion and (diluted) collyrium.

Incomp. Acids and alkalies.

FERRUM.**1. FERRI SESQUIOXYDUM. (Ferri oxydum, F. carbonas, F. subcarbonas, F. rubigo.)**

Dose and Form. Gr. x-3 ij, as an electuary with syrup, treacle, and with aromatics.

Incomp. Acids and acidulous salts.

Use. As a general tonic, especially in anaemic condition.

2. FERRI AMMONIO-CHLORIDUM. (F. ammonium.)

Dose. Gr. iij– $\frac{3}{4}$ j in powder, or pill with extracts.
 (Little used.)

3. TINCTURA FERRI AMMONIO-CHLORIDI.

Comp. F. Am. chlor. $\frac{3}{4}$ iv, proof spir. f $\frac{7}{3}$ xx.

Dose. $\frac{3}{4}$ ss-ij. (Little used.)

4. TINCTURA FERRI SESQUICHLORIDI.

(Tinct. Ferri muriatis.)

Comp. F. subcarb. flss, acid. hydrochlor. Oj,
 alc. Oijj.

Dose. $\frac{3}{4}$ v-x- $\frac{3}{4}$ j, in simple liquids.

Incomp. Alkalies and their carbonates, astringent
 vegetable infusions, gum Arabic.

Use. A general tonic with special tendency to-
 wards the urinary organs.

5. FERRI IODIDUM.

Comp. A deliquescent salt : iron 14, iodine 63·3,
 water 22·7 in 100 parts.

Dose. Gr. ij-vj, in distilled water with tinct. (with
 iron wire.)

Incomp. Acids, alkalies, vegetable infusions, &c.
 (Should always be kept ready in solution, else it
 is never neutral.)

(See DEOBSTRUENTS OR ALTERATIVES.)

6. FERRI PERCYANIDUM. (Hardly used medici-
 nally.)

7. POTASSII FERROCYANIDUM. (Do.)

8. FERRI SULPHAS.

Dose. Gr. j-v, in pill, with extracts or aromatic
 confection.

9. MISTURA FERRI COMPOSITA.

(Griffith's mixture.)

Comp. Myrrh 3 j, carb. of potash gr. xxv, sulphate of iron 3 j, sugar 3 j, spirit of lavender 3 ss, rose water 3 vijss.

Dose. 3 ss-ij, simple or with aromatics.

Incomp. Acids, acidulous salts, bitters.

Use. In chlorosis, menstrual derangement, hectic.

10. PILULÆ FERRI COMPOSITÆ.

Comp. 1 sulphate of iron, 1 carbonate of soda, syr. q. s., 2 myrrh. (The sulphate is decomposed.)

Use. A good emmenagogue.

11. FERRI POTASSIO-TARTRAS. (F. Tartarisatum.)

Dose. Gr. x-3 ss in pill, bolus, or solution in carminative water.

ZINCUM.

Use. For cases of debility attended with irritation.

1. ZINCI OXYDUM. (Flores Zinci.)

Dose. Gr. j-vj, in pill or powder.

2. ZINCI SULPHAS.

Dose. Gr. j-v, in pill.

VEGETABLE TONICS. BITTER TONICS.

CALUMBA. (Cocculus Palmatus. Root.)

Dose. Gr. x-3 ; in powder.

Use. In decline of remittent, of dysentery, of cholera.

1. INFUSUM CALUMBÆ.

Comp. Calumba 3 ss, water Oj.

Dose. 3 j-ij.

Incomp. Bichloride of mercury, lime water.

2. TINCTURA CALUMBÆ.

Comp. Calumba $\frac{3}{2}$ iv, dil. alc. Oij.

Dose. f $\frac{3}{2}$ j-iv (added to bitter infusions and effervescent draughts).

CENTAURIUM. (Erythræa C. The flowers and leaves.)

Dose and Form. Of the powder, $\frac{3}{2}$ j-3 j. Infusion ($\frac{3}{2}$ ij- $\frac{3}{2}$ x water) $\frac{3}{2}$ i-ij.

Use. In debilitated states of the stomach.

CINCHONA. (a. C. Cordifolia, Yellow bark.
b. C. Lancifolia, Pale bark. c. C. Oblongifolia, Red bark.)

Dose. Powder, gr. x- $\frac{3}{2}$ ij.

Use. Intermittent fever, whenever a permanent corroborative effect is desired, provided the stomach be in a proper state.

1. INFUSUM CINCHONÆ.

Comp. Cinchona (lancifol.) $\frac{3}{2}$ j, water Oi.

Dose. f $\frac{3}{2}$ j-ij.

Incomp. Lime water, tart. emet., sulphate of zinc and iron.

**2. DECOCTUM CINCHONÆ CORDIFOLIÆ. (Olim
DECOCTUM CINCHONÆ.)****3. DECOCTUM CINCHONÆ LANCIFOLIÆ.****4. DECOCTUM CINCHONÆ OBLONGIFOLIÆ.**

Comp. Cinchona $\frac{3}{2}$ j, water Oj.

Dose. $\frac{3}{2}$ j-ij.

Incomp. Lime water, tartar emetic, sulphate of iron.

5. TINCTURA CINCHONÆ.

Comp. C. cord. $\frac{2}{3}$ vj, dil. alc. Oij.

Dose. f $\frac{3}{4}$ j- $\frac{2}{3}$ ss.

6. TINCTURA CINCHONÆ COMPOSITA. (Huxham's Tincture.)

Comp. Cinchona (lanc.) $\frac{2}{3}$ ij, orange peel $\frac{2}{3}$ iss, serpentary $\frac{3}{4}$ iij, saffron $\frac{3}{4}$ j, red saunders $\frac{3}{4}$ j, dil. alc. f $\frac{2}{3}$ xx.

Dose. f $\frac{3}{4}$ j-iv.

7. EXTRACTUM CINCHONÆ CORDIFOLIÆ. (Olim EXTRACT. CINCHONÆ.)

8. EXTRACTUM CINCHONÆ LANCIFOLIÆ.

9. EXTRACTUM CINCHONÆ OBLONGIFOLIÆ.

Comp. Watery extract.

Dose. Gr. x-xx.

Form. Pill, solution in water, &c., syrup of mulberries, (for children.)

10. QUINÆ DISULPHAS. (Sulphate of quinine.)

Comp. From the C. cordifolia.

Dose. Gr. ss-x.

Form. Pill, with conserve of roses, powder with sugar. Solution, in acidulated liquids, water, alcohol or syrup.

[The best vehicle is distilled water, slightly acidulated. It is little soluble in pure water; more so in spirit. Soluble in 780 parts of cold water, or 80 parts of spirit of sp. gr. 0.85. (Perreira.) Very soluble in sulph. acid. A drachm of acid. sulph. dil. will dissolve gr. xx. A convenient strong solution is made by dissolving gr. j. in water $\frac{3}{4}$ j, by the addition of acid. sulph. dil. $\frac{1}{2}$ jj. When infusion of roses is

used as a vehicle, dilute nitric acid should be used in place of sulphuric, and in the proportion of *half as much more* as of the sulphuric. The nitric acid keeps the tannate of quinine in solution, and keeps the mixture clear. (Mr. Squire, in Pharm. Jour.)]

Use. Preferable to cinchona from convenience of administration. Possesses more anti-intermittent and less tonic property than cinchona.

GENTIANA. (G. Lutea. The root.)

Dose. Powder gr. x-3 ss. (Not much used.)

Use. In dyspepsia, gout, worms, intermittent.

1. INFUSUM GENTIANÆ COMPOSITUM.

Comp. Gentian $\frac{3}{2}$ ss, orange peel 3 i, coriander 3 j, dil. alc. $\frac{3}{2}$ iv, water $\frac{3}{2}$ xij.

Dose. $\frac{3}{2}$ j-ij.

Incomp. Sulphate of iron and zinc.

2. MISTURA GENTIANÆ COMPOSITA.

Comp. Inf. gent. comp. f $\frac{3}{2}$ vj, inf. sennæ comp. f $\frac{3}{2}$ iij, tinct. card. c. f $\frac{3}{2}$ j.

Dose. f $\frac{3}{2}$ x- $\frac{3}{2}$ ij.

3. TINCTURA GENTIANÆ COMPOSITÆ.

Comp. Gentian $\frac{3}{2}$ ii, orange peel $\frac{3}{2}$ j, cardam. $\frac{3}{2}$ ss, dil. alc. Oij.

Dose. f $\frac{3}{2}$ ss-ij.

4. EXTRACTUM GENTIANÆ. (Watery extract.)

Dose. Gr. x-3 ss.

MENYANTHES. (M. trifoliata. Buckbean. The leaves.)

Form and Dose. Powder $\frac{3}{2}$ j-3 ss; infusion ($\frac{3}{2}$ j-to $\frac{3}{2}$ xx of water) $\frac{3}{2}$ j-iss; extract (not much used).

Use. Unites a cathartic with tonic property.

QUASSIA. (*Q. excelsa.* The wood.)

Dose and Form. Gr. x-3 ss of the powder.
(Rarely used.)

1. INFUSUM QUASSIÆ.

Comp. Quassia 3 ij, cold water Oj.

Dose. f 3 j-ij.

Incomp. Lead, silver. (May be used with chalybeates.)

Use. A pure bitter. In dyspepsia from debility of stomach, in remission of fever.

SIMARUBA. (*S. officinalis.* The bark of the root.)

Dose. 3 ss-3 ss in powder. (Rarely used.)

Use. Mostly in Europe. Dysentery.

1. INFUSUM SIMARUBÆ.

Comp. Simaruba 3 iiij, water 3 xx.

Dose. 3 j-ij.

Incomp. Alkalies and their carbonates, lime water.

PART II.

REMEDIES FOR EXTERNAL AND
LOCAL USE.*Cataplasma. Poultices.*

EMOLlient. SEDATIVE.

1. CATAPlASMA LINI. Linseed-meal poultice.
Comp. Boiling water f $\frac{3}{2}$ xx, linseed meal q. s.
2. CATAPlASMA CONII. Hemlock poultice.
Comp. Boiling water $\frac{3}{2}$ xx, extract. conii $\frac{3}{2}$ ij,
linseed meal q. s.
3. [CATAPlASMA ALUMINIS. Ph. Dub. Alum poultice.
Comp. White of two eggs, alum 3 j.]

STIMULANT. ANTISEPTIC.

1. CATAPlASMA SINAPIS. Mustard poultice.
Comp. Powdered mustard and linseed meal, of each $\frac{3}{2}$ v, boiling vinegar q. s.
2. ALLIUM. Garlic. Boiled to a proper consistency.
3. CATAPlASMA FERMENTI. Yeast poultice.
Comp. Flour $\frac{3}{2}$ xij, yeast of beer $\frac{3}{2}$ x.

4. [CATAPLASMA CARBONIS LINI. Ph. Dub. Charcoal poultice.
Comp. Charcoal q. s., added to linseed or oatmeal poultice.]
 5. [CATAPLASMA DAUCI. Carrot poultice. Ph. Dub. The carrot boiled to a proper consistence.]
-

Caustica. Vesicantia. Irritantia.

(Caustics. Blisters. Counter-irritants.)

1. POTASSÆ HYDRAS. (Olim P. Fusa.)
2. POTASSA CUM CALCE. Less deliquescent than the preceding.
3. ARGENTI NITRAS.
4. ACIDUM ACETICUM.
5. EMPLASTRUM CANTHARIDIS.
Comp. 1 lard, 2 cantharides, 3 emplast. ceræ.
6. ACETUM CANTHARIDIS.
Comp. Cantharides $\frac{3}{4}$ j, acid. acet. $\frac{3}{4}$ x.
7. [Tela Vesicatoria. Charta Vesicatoria. More elegant and equally effective preparations. Cantharadin, &c.]
8. CERATUM CANTHARIDIS.
Comp. Powdered cantharides $\frac{1}{2}$ j, yellow wax and lard each $\frac{3}{4}$ viij.
9. UNGUENTUM CANTHARIDIS.
Comp. 1 decoct. canthar., 2 cerat. resinae.

10. CERATUM SABINÆ.

Comp. Savine $\frac{3}{4}$ ij, resin cerate lbj.

11. [Unguentum Mezerei. Amer. Ph.

Comp. Mezereon $\frac{3}{4}$ iv, lard $\frac{3}{4}$ xiv, wax $\frac{3}{4}$ ij.]

12. UNGUENTUM ANTIMONII POTASSIO-TARTRATIS.

Tartar-emetic ointment.

Comp. Ant. pot. tart. $\frac{3}{4}$ ij, lard $\frac{3}{4}$ j.

13. EUPHORBIUM. Mixed with turpentine or Bur-gundy pitch.

14. CAPSICUM. Added to Cataplasms.

15. MEZEREON. The bark softened in hot vinegar.

16. PIPER NIGRUM. Mixed with sinapisms or cataplasms.

17. RUTA. In substance, as a cataplasm.

[MOXAS.]

Substances for producing counter-irritation by being burnt on the part. They are composed of various materials, and may be composed of any that will burn down slowly, such as cotton, the pith of plants, agaric, German tinder, &c. Percy used an *artillery match*. Larrey's moxa was a truncated cone of cotton stitched in linen, about four lines wide and six lines high. Dr. Sadler's, of St. Petersburg, are composed of the pith of the sunflower, cotton, and linen, steeped in a solution of nitre. Gräfe employed wafers dipped in three parts of oil of turpentine and one part of æther.]

Cerata et Unguenta. Ointments.

EMOLlient. DEFENSIVE.

1. CERATUM.

Comp. Wax and olive oil, equal parts.

2. CERATUM CALAMINÆ. (Turner's cerate.)

Comp. Calamine, wax, $\frac{1}{2}$ ss, lard $\frac{1}{2}$ ij.

3. CERATUM CETACEI. Spermaceti cerate.

Comp. Cetac. $\frac{1}{2}$ j, ceræ $\frac{1}{2}$ iiij, ol. oliv. f $\frac{1}{2}$ vj.

4. CERATUM PLUMBI ACETATIS.

Comp. Plumb. acet. $\frac{1}{2}$ ij, ceræ $\frac{1}{2}$ ij, ol. oliv. $\frac{1}{2}$ viij.

5. CERATUM PLUMBI COMPOSITUM. (Goulard's cerate.)

Comp. Liq. plumb. diacet. f $\frac{1}{2}$ iiiss, ceræ $\frac{1}{2}$ iv, ol. oliv. f $\frac{1}{2}$ ix, camphoræ $\frac{1}{2}$ ss.

6. CERATUM SAPONIS.

Comp. Sol. subacetate of lead Oij, wax $\frac{1}{2}$ x, olive oil Oj, soap $\frac{1}{2}$ vj.

7. UNGUENTUM CETACEI.

Comp. Spermaceti $\frac{1}{2}$ vj, wax $\frac{1}{2}$ ij, ol. oliv. $\frac{1}{2}$ iiij.

8. UNGUENTUM SAMBUCI. (Elder ointment.)

Comp. Elder flowers and lard, equal parts—boil and strain.

9. UNGUENTUM ZINCI.

Comp. 1 oxide of zinc, 6 lard.

STIMULANT. DIGESTIVE.

1. CERATUM RESINÆ. (Basilicon.)

Comp. Resin $\frac{1}{2}$ v, wax $\frac{1}{2}$ ij, lard $\frac{1}{2}$ viij,

2. UNGUENTUM CREOSOTI.

Comp. Creosote $\frac{1}{2}$ ss, lard $\frac{1}{2}$ j.

3. UNGUENTUM ELEMI.

Comp. Elem. $\frac{3}{2}$ vj, turpentine $\frac{3}{2}$ v, suet $\frac{3}{2}$ xij, ol. oliv. f $\frac{3}{2}$ ij.

4. UNGUENTUM GALLÆ COMPOSITUM.

Comp. Powdered galls $\frac{3}{2}$ ij, lard $\frac{3}{2}$ ij, opium $\frac{3}{2}$ ss.

5. UNGUENTUM HYDRARGYRI AMMONIO-CHLORIDI. (White precipitate ointment.)

Comp. Hydr. am. chlor. (white precipitate) $\frac{3}{2}$ j, ung. simp. $\frac{3}{2}$ iss.

6. UNGUENTUM HYDRARGYRI NITRATIS. (Citrine ointment.)

Comp. Mercury $\frac{3}{2}$ j, ac. nit. $\frac{3}{2}$ xij, fresh neat's foot oil f $\frac{3}{2}$ ix, lard $\frac{3}{2}$ iij.

7. UNGUENTUM HYDRARGYRI NITRICO-OXIDI. (Red precipitate ointment.)

Comp. 1 hyd. nit. ox., 8 ung. simp.

8. UNGUENTUM PICIS LIQUIDI. (Tar ointment.)

Comp. Suet and tar, equal parts.

9. UNGUENTUM PICIS NIGRI. (Pitch ointment.)

Comp. Pitch, wax, resin $\ddot{\text{aa}}$ $\frac{3}{2}$ ix, ol. oil $\frac{3}{2}$ xvij.

10. UNGUENTUM PLUMBI COMPOSITUM.

Comp. Cret. præp. $\frac{3}{2}$ iv, acet. destil. $\frac{3}{2}$ iij, emplast. plumb. $\frac{3}{2}$ xvij, ol. oliv. $\frac{3}{2}$ x.

ALTERATIVE. DISCUTIENT.

1. CERATUM HYDRARGYRI COMPOSITUM.

Comp. 4 unguent. hyd. fort., 4 cerat. sap., 1 camphor.

2. UNGUENTUM HYDRARGYRI FORTIUS.

Comp. 24 mercury, 23 lard, 1 suet.

3. UNGUENTUM HYDRARGYRI MITIUS.

Comp. 1 unguent. hyd. fort., 2 lard.

4. UNGUENTUM HYDRARGYRI IODIDI.

Comp. 1 iod. of merc., 2 wax, 6 lard,

5. UNGUENTUM HYDRARGYRI BINIODIDI.

Comp. 1 biniod. of merc., 2 wax, 6 lard. (More active than the preceding.)

6. UNGUENTUM IODINII.

Comp. Iodine 3 j, lard 3 j, alc. ℥ xx.

7. UNGUENTUM IODINII COMPOSITUM.

Comp. Iodine 3 ss, iod. of potass. 3 j, lard 3 ij alc. 3 j.

8. UNGUENTUM PLUMBI IODIDI.

Comp. 1 iodide of lead, 8 lard.

SPECIFIC.

1. UNGUENTUM SULPHURIS.

Comp. Sulphur ℥ jj, lard ℥ jjij.

2. UNGUENTUM SULPHURIS COMPOSITUM.

Comp. Sulphur 3 j, ammon. merc. et acid benzoinic aa 3 j, ol. bergamot et acid sulph. aa f 3 j, nit. pot. 3 ij, lard ℥ ss.

3. UNGUENTUM VERATRI.

Comp. White hellebore 3 ij, lard 3 viij, ol. limon. ℥ xx.

SEDATIVE. ANTISPASMODIC.

[1. *Unguentum Veratriæ.* Veratria 3 j, olive oil 3 j, lard 3 vij.

2. *Unguentum Aconitinæ.* Aconit. gr. iv, ol. oliv. 3 ss, lard 3 j.

3. *Unguentum Belladonnæ.* Extr. Bell. 3 j-ij, lar l, or spermaceti ointment 3 j.]

Emplastra. Plasters.

STIMULANT. DISCUTIENT.

1. EMPLASTRUM AMMONIACI.

Comp. Ammoniacum $\frac{3}{2}$ v, distil. vinegar Oss.

2. EMPLASTRUM AMMONIACI CUM HYDRARGYRO.

Comp. Ammoniac. $\frac{3}{2}$ xij, mercury $\frac{3}{2}$ iij, oil and sulphur q. s.

3. EMPLASTRUM GALBANI.

Comp. Galbanum $\frac{3}{2}$ ij, emplast. plumb. $\frac{3}{2}$ ix, turpentine $\frac{3}{2}$ iiss, resin $\frac{3}{2}$ vj.

4. EMPLASTRUM HYDRARGYRI.

Comp. Mercury $\frac{3}{2}$ vj, emplast. plumb. $\frac{1}{2}$ vj, olive oil and resin, each $\frac{3}{2}$ ij.

ANODYNE. EMOLLIENT.

1. EMPLASTRUM BELLADONNÆ.

Comp. Extr. belladonnæ $\frac{3}{2}$ iss, emplast. resinæ $\frac{3}{2}$ iij.

2. EMPLASTRUM OPII.

Comp. Opium $\frac{3}{2}$ ij, pic. Burgund. $\frac{3}{2}$ iij, emplast. plumb. $\frac{1}{2}$ vj, water $\frac{3}{2}$ iv.

ADHESIVE. DEFENSIVE.

1. EMPLASTRUM CERÆ.

Comp. 1 resin, 3 wax, 3 suet.

2. EMPLASTRUM PLUMBI.

Comp. Semivitrif. ox. plumb. $\frac{1}{2}$ v, olive oil Cong. j, water Oij.

3. EMPLASTRUM SAPONIS.

Comp. Soap $\frac{1}{2}$ ss, emplast. plumbi. $\frac{1}{2}$ iiij.

4. [Court plaster. Solution of isinglass on silk, and tincture of benzoin.]
 5. Liston's plaster. A solution of isinglass on oiled silk.]
-

Enemata. Alvine Injections.

CATHARTIC. STIMULANT.

1. ENEMA ALÖES.

Comp. Aloes 3 ij, carb. potass. gr. xv, decoct. hordei 3 x.

2. ENEMA COLOCYNTHIDIS.

Comp. Extract. coloc. c. 3 ij, soap 3 j, water 3 xx.

3. ENEMA TEREBINTHINÆ.

Comp. Ol. terebinth. 3 j, yolk of eggs, q. s., hot water 3 xix.

4. [ENEMA CATHARTICUM. Ph. Dub.

Comp. Manna 3 j, magnes. sulph. 3 ss, ol. oliv. 3 j, decoct. anthem. 3 x.]

5. [ENEMA RUTÆ.

Comp. Confect. Rutæ 3 j, decoct. hord. 3 vj.]

SEDATIVE. EMOLLIENT.

1. ENEMA OPII.

Comp. Tinct. opii 3 ss, decoct. of starch 3 iv.

2. ENEMA TABACI.

Comp. Tobacco 3 j, boiling water 3 xx.

3. [ENEMA FÆTIDUM Ed. Dub.

Comp. Enema cathart. 3 xij, tinct. assaf. 3 ij.]

4. [ENEMA EMOLLIENS. (Decoct. malvæ comp.)

Comp. Mallow 3 j, chamomile 3 ss, water 3 xx, —boil and strain.]

The Endermic Method.

This consists in applying medicines in fine powder, pure or mixed into a paste with lard, gelatine, or cerate, to a blistered surface. The following medicines, among many others, have been used in this manner, and in the doses mentioned :

Morphia, and its salts	gr. $\frac{1}{4}$ -ij.
Strychnia	gr. $\frac{1}{4}$ -j.
Aconitina	gr. $\frac{1}{16}$ - $\frac{1}{8}$.
Extract. belladonnæ	gr. ij-v.
Quinæ disulphas	gr. ij-vj.
Moschus	gr. iv-viij.

Gargarismata. Gargles.

USED UNDILUTED.

1. DECOCTUM CINCHONÆ.
2. DECOCTUM QUERCUS.
3. INFUSUM CATECHU COMPOSITUM.
4. INFUSUM CINCHONÆ.
5. INFUSUM ROSÆ COMPOSITUM.

USED DILUTED.

6. LINIMENTUM ÆRUGINIS.
7. MEL BORACIS.
8. TINCTURA GALLÆ.
9. TINCTURA CAPSICI.
10. TINCTURA KINO.
11. TINCTURA MYRRHÆ.

Linimenta. Embrocationes. Fomentationes. Lotiones.

STIMULANT.

1. LINIMENTUM AMMONIÆ. (Stronger liniment.)

Comp. 1 liquor ammoniæ, 2 olive oil.

2. LINIMENTUM AMMONIÆ SESQUICARBONATIS.
(Weaker liniment.)

Comp. 1 liq. amm. sesquicarb., 3 olive oil.

3. [LINIMENTUM AMMONIÆ COMPOSITUM.

Comp. Liq. ammoniæ $\frac{2}{3}$ v, tinct. camph. $\frac{2}{3}$ ij,
spir. rosmar. $\frac{2}{3}$ j. Ph. Ed.]

[The preceding formula is an imitation of Dr. Granville's *antidynous lotions*, which are,

a. *Lotio Ammoniata mitior.*

Comp. Liq. ammon. fortiss. $\frac{2}{3}$ j, spir. rosmar.
 $\frac{2}{3}$ vj, spir. camph. $\frac{2}{3}$ ij.

b. *Lotio Ammoniata fortior.*

Comp. Liq. ammon. fortiss. $\frac{2}{3}$ x, spir. rosmar.
 $\frac{2}{3}$ ss, spir. camph. $\frac{2}{3}$ ij.]

4. LINIMENTUM CAMPHORÆ.

Comp. Camphor $\frac{2}{3}$ ss, olive oil $\frac{2}{3}$ ij.

5. LINIMENTUM CAMPHORÆ COMPOSITUM.

Comp. Camphor $\frac{2}{3}$ iiiss, liq. ammoniæ $\frac{2}{3}$ viiss,
spir. lavand. $\frac{2}{3}$ xx.

6. LINIMENTUM HYDRARGYRI COMPOSITUM.

Comp. Unguent. hyd. fort. $\frac{2}{3}$ iv, lard $\frac{2}{3}$ iv, camphor $\frac{2}{3}$ j, liq. ammon. $\frac{2}{3}$ iv.

7. LINIMENTUM SAPONIS.

Comp. Soap $\frac{2}{3}$ iij, camphor $\frac{2}{3}$ j, oil rosmar. f $\frac{2}{3}$ j,
ol. origan. f $\frac{2}{3}$ j, alcohol Oj.

8. LINIMENTUM TEREBINTHINÆ.

Comp. Oil of turpentine Oss, cerat. resin ℥vj.

9. OLEUM CAJAPUTI.

Comp. Simple, or mixed with olive oil.

10. [LINIMENTUM CROTONIS.]

Comp. 1 croton oil, 5 olive oil.]

REFRIGERANT.

1. LIQUOR PLUMBI DIACETATIS. (Goulard.)

2. LIQUOR PLUMBI DIACETATIS DILUTUS.

Comp. Liq. pl. diacet. f 3 ij, dist. water Oj.

3. [Lotio Ammoniae Hydrochloratis. (Muriate of ammonia wash.)]

Comp. Ammon. hydrochlor. ʒ j-ij, aquæ vel aceti ʒ xij, spir. rect. ʒ iv.]

SEDATIVE. EMOLlient.

1. DECOCTUM PAPAVERIS.

Comp. Poppy heads ʒ j, water f ʒ xx,—boil 15 minutes.

2. LINIMENTUM OPII.

Comp. 1 tinct. opii, 3 liniment. saponis.

3. DECOCTUM MALVÆ COMPOSITUM.

Comp. Mallow ʒ j, chamomile ʒ ss, water f ʒ xx.

[*Embrocatio Aconitinæ.* Aconit. gr. viij, spir. rect. ʒ ij.

Embrocatio Aconiti. Extract. Aconiti ʒ ss ad ʒ j, alcohol ʒ ij tere bene et cola. This answers well, and is only one-sixth part the price of the preceding].

ASTRINGENT.

1. LIQUOR ALUMINIS COMPOSITUS.

Comp. Alum 3 ij, sulph. zinci 3 ij, water f 3 xv.

2. LIQUOR CALCIS.

Comp. Lime 3 iv, dist. water Cong. j.

3. DECOCTUM QUERCUS.

Comp. Oak bark 3 j, water Oiss—boil to Oj.

FOR WOUNDS AND SKIN DISEASES.

1. LINIMENTUM ÆRUGINIS.

Comp. 1 verdigris, 7 vinegar, 14 honey.

2. LIQUOR CUPRI AMMONIO-SULPHATIS.

Comp. Cupri am. sulph. 3 ss, water 3 x.

3. DECOCTUM VERATRI.

Comp. Verat. alb. 3 x, water 3 xl, (boil to 3 xx), spir. rect. 3 iij.

4. [*Linimentum Calcis.* (Carron oil.)

Comp. Equal parts of lime water and linseed oil.]

5. [*Lotio Flava.* (Aqua Phagedenica.)

Comp. Hydrarg. bichlor. gr. ij, aquæ calcis 3 j.]

6. [*Lotio Nigra.* (Black wash.)

Comp. Calomel 3 j, liq. calcis 3 vj.]

7. [*Lotio Hydrocyanica.* Acid. hydroc. dilut. 3 j, water 3 ij.]

Pulveres. Powders.

DESICCANT. ASTRINGENT. STIMULANT.

1. CALAMINA PRÆPARATA.
 2. CREA PRÆPARATA.
 3. ALUMEN EXSICCATUM.
 4. RHEI PULVIS.
 5. HYDRARGYRI NITRICO-OXIDUM.
 6. HYDRARGYRI CHLORIDUM.
-

Inhalation.

Medicines are applied to the mucous membrane of the larynx, trachea, and bronchi, in several ways.

1. Diffused in the air of the chamber, and thus inhaled in ordinary respiration. The vapour of water (steam), the fumes of tar, resin, myrrh, &c., are thus applied.

2. Mixed with water in a close glass vessel having two tubes inserted into it, the one admitting the air and passing it through the mixture, the other inserted into the patient's mouth, through which the air of the vessel is drawn into the lungs. Chlorine gas, tincture of iodine, of conium, &c., are thus inhaled.

3. By means of the apparatus called "Mudge's Inhaler," or a greatly improved modification of this by Dr. Harwood, having a mouthpiece, which renders the process of inhalation much easier. Warm, watery vapours, in which various medicaments may be diffused or developed, are thus applied to the air-passages.

PART III.

I. POISONS AND COUNTERPOISONS.

Corrosive and Irritant Poisons.

POISONS.

Arsenic.

The strong Mineral Acids.
Nitric A.
Muriatic A.
Sulphuric A.

TREATMENT.*

(Sulphas Zinci. Sinapis Pulvis.
Ipecacuanha.)

The stomach-pump. Emetics.
Milk in copious draughts. Oil
and lime water in equal parts.
Mucilage. Flour and water. Hy-
drated oxide of iron, recently pre-
pared. (*Ferri sesquioxidum* Lond.
Carbonas Ferri præcipitatis Ed.
Ferri carbonas Dub.)

Magnesia, chalk, whiting, in
milk or water, mucilaginous or
soapy liquid in large quantities.

* In this table the primary or immediate treat-
ment is alone, in general, noticed.

POISONS.

TREATMENT.

Oxalic Acid.

Stomach-pump. Mechanical irritation of the fauces. Magnesia, chalk, whiting, in small quantities of liquid. Muriate of lime diluted.

The strong or Caustic Alkalies.

Ammonia.

Potassa.

Soda.

Lime.

Stomach-pump. Emetics, Diluents. Demulcents. Olive oil. Diluted vinegar, lemon-juice, or citric acid, ad libitum.

*Antimony,
Salts of.*

Tartar Emetic.

Butter of Antimony.

Copious draughts of warm water. Watery infusions containing gallic acid. Infusion of galls, oak bark, gentian, cascara, calumba, cinchona flava, kino, catechu.

Baryta.

Barii Chloridum.

Alkaline sulphates. Sulphate of soda or magnesia. Stomach-pump.

Carbonas Barytæ.

Sulphate of soda or magnesia mixed with acetic acid.

Copper, Salts of.

Acetate.

Carbonate.

Nitrate.

Sulphate.

Demulcents. Albumen. Raw eggs. Flour and water. Milk, cream.

Iodine.

Emetics. Stomach-pump. Tepid demulcents. Flour, sago, arrow-root?

POISONS.

TREATMENT.

<i>Lead, Salts of.</i>	Emetics. Stomach-pump. Alkaline sulphates, especially S. magnesiae.
<i>Carbonate.</i>	Sulphate of soda mixed with acetic acid.
<i>Mercury.</i> <i>Salts of.</i> Bichloride (corrosive S.)	Albumen, or beaten eggs in large quantity. Flour and water in stiff paste (gluten). Milk.
Bicyanide. Iodide. Nitro-oxide.	
<i>Nitre.</i> <i>Phosphorus.</i>	Warm demulcents, Stomach-pump. Cold water. Magnesia, &c. as for the mineral acids.
<i>Silver, Salts of.</i> Nitrate (lunar caustic).	Chloride of sodium (common salt) in large quantity of water. Stomach-pump.
<i>Tin, Salts of.</i> Chloride (dyer's liquid).	Stomach-pump. Emetics, Diluents. Demulcents. Milk, &c.
<i>Zinc, Salts of.</i> Sulphate.	Demulcents. Diluents. Milk. Albumen.

Narcotic and Narcotico-Acid Poisons.

POISONS.	TREATMENT.
<i>Gaseous Poisons</i>	The open air. Semi-erect position. Cold water to the face and neck, if the surface is warm.—Mechanical excitement of respiration by manipulating the thorax and abdomen. Feet in hot water. V. S. cucurb. nuchæ. Flagellation. Lin. camph. comp. to chest.
Carbonic acid.	
Sulphuretted Hydrogen.	
Carburetted Hydrogen.	
<i>Opium.</i>	Stomach-pump. Emetics. Mechanical irritation of the fauces. External excitation. Blows on the palms and soles. Semi-erect position. Mechanical excitement of respiration. Cold affusion to the head and face. Feet in hot water. V. S. cucurb. nuchæ. Internal stimulants. Ammonia. Coffee. Vegetable acids.
Morphia, &c.	
<i>Hydrocyanic Acid.</i>	Same as in opium. Inhalation of vapour of ammonia much diluted. Cold affusion. Mixed oxides of iron, obtained by precipitating solution of green sulphate with potash.
<i>Oil of Bitter Almonds.</i>	Same as in hydrocyanic acid.
<i>Oil and Water of Cherry-laurel.</i>	The same.

POISONS.

TREATMENT.

Aconitum Nau-

pellus (Aconite).

Agaricus.

Alcohol (Coma-

tose Intoxica-

tion).

Atropa Bellata-

donna,

(Deadly

Nightshade).

Cicuta Virosa,

(Water Hem-

lock).

Conium Macu-

latum,

(Spotted

Hemlock).

Œnanthe Apfo-

lia.

Œnanthe Cro-

cata.

Cocculus Indi-

cus.

Colchicum.

Digitalis.

Ergota.

Helleborus Ni-

ger, (Black

Hellebore).

Nux Vomica.

Stramonium.

Strychnine.

Tobacco.

Veratrum Al-

bum, (White

Hellebore).

The general treatment in poison-ing by the other simple narcotics and acrid narcotics in this list, is the same as in cases of opium and hydrocyanic acid ; the chief differences depending on the amount of local irritation produced, and the consequent administration of refrigerants, mild opiates, &c. Emetics, and stomach-pump.

(In poisoning from *digitalis*, the recumbent posture to prevent syn-cope.)

Acrid Poisons.

POISONS.	TREATMENT.
<i>Brionia Dioica</i> , (Wild Vine).	Vomiting by ipecacuanha and irritation of the fauces. Copious use of diluents and demulcents.
<i>Cantharides</i> .	
<i>Elaterium</i> .	Opiates to relieve pain, mild cor- dials, &c.
<i>Euphorbium</i> .	

II. TEMPERATURE OF BATHS.

	WATER.	VAPOUR.	
		Not breathed.	Breathed.
Cold Bath	33°—63°		
Cool Bath	63°—75°		
Temperate Bath...	75°—85°		
Tepid Bath	85°—92°	96°—106°	90°—100°
Warm Bath.....	92°—98°	106°—120°	100°—110°
Hot Bath	98°—106°	120°—160°	110°—130°

III. QUANTITY OF SOME OF THE STRONGER
MEDICINES IN CERTAIN PHARMACEUTI-
CAL COMPOUNDS.

1. OPIUM.

Confectio opii contains gr. j in gr. xxxvj.

Enema opii, gr. iss. in $\frac{3}{4}$ iv.

Linimentum opii, gr. iij in $\frac{3}{4}$ ss.

Pilulæ saponis compositæ, gr. j in gr. v.

Pilulæ styracis compositæ, gr. j in gr. v.

Pulvis cretæ comp. cum opio, gr. j in $\frac{3}{4}$ ij.

Pulvis ipecacuanhæ compositus, gr. j in gr. x.

Pulvis kino compositus, gr. j in $\frac{3}{4}$ j.

Tinctura camphoræ comp., gr. j in $\frac{3}{4}$ ss.

Tinctura opii, gr. j in $\frac{3}{4}$ xij.

Vinum opii, gr. j in $\frac{3}{4}$ xij.

2. MERCURY.

Hydrargyrum cum creta, gr. iij in gr. viij.

Linimentum hydrargyri comp., $\frac{3}{4}$ j to $\frac{3}{4}$ xj.

Liquor hydrargyri bichloridi, gr. j (of hyd. bichlor.)
in $\frac{3}{4}$ xij.

Pilulæ hydrargyri, gr. j in gr. iiij.

Pilulæ hydrarg. chloridi comp., gr. j (of calomel) in
gr. v.

Pilulæ hydrargyri iodidi, gr. j (of hydrarg. iodid.)
in gr. v.

Unguentum hydrargyri fortius, $\frac{3}{4}$ j in $\frac{3}{4}$ ij.

Unguentum hydrargyri mitius, $\frac{3}{4}$ j in $\frac{3}{4}$ vj.

3. ARSENIC.

Liquor potassæ arsenitis, gr. j in $\frac{2}{3}$ iiss.

4. ANTIMONY.

Vin. ant. pot. tart., gr. j in $\frac{2}{3}$ ss.

Unguentum ant. pot. tart., $\frac{3}{4}$ j in $\frac{2}{3}$ ss.

IV. PROPORTION OF DOSES ACCORDING TO
AGE. (GAUBIUS.)

For an adult,

Suppose the dose to be ONE or 1 drachm.

Under 1 year	will only require	. . .	$\frac{1}{12}$	5 grains.
2		. . .	$\frac{1}{8}$	8 grains.
3		. . .	$\frac{1}{6}$	10 grains.
4		. . .	$\frac{1}{4}$	15 grains.
7		. . .	$\frac{1}{3}$	1 scruple.
14		. . .	$\frac{1}{2}$	$\frac{1}{2}$ drachm.
20		. . .	$\frac{2}{3}$	2 scruples.
21 to 60	the full dose ONE			1 drachm.

Above this age an inverse gradation must be observed.

V. CLASSIFIED LIST OF THE PRINCIPAL
MINERAL WATERS.

I. Saline aperient.

1. Carlsbad (165°).
2. Kissingen (cold).
3. Marienbad (cold).
4. Franzensbad (cold).
5. Cheltenham (cold).
6. Leamington (cold).
7. Scarborough (cold).

II. Hot saline.

1. Wiesbaden (160°).
2. Baden-Baden (153°).

III. Hot sulphurous.

1. Aix-la-Chapelle (130°).
2. Barèges (130°).
3. Bagnères de Luchon (130°).

IV. Hot alkaline.

1. Vichy (108°).
2. Mont d'Or.
3. Ems (83°-115°).
4. Töplitz (114°-122°).
5. Schlangenbad (86°).
6. Wildbad (88°-99°).

*v. Simple unmineralized,
hot.*

1. Pfessers (100°).
2. Gastein (118°).
3. Buxton (82°).
4. Bristol (74°).
5. Matlock (66°).

VI. Simple mineralized, hot.

1. Bath (112°-116°).
2. Pyrenees (80°-122°).

VII. Chalybeate waters.

1. Spa (cold).
2. Schwalbach (cold).
3. Pyrmont (cold).
4. Marienbad (cold).
5. Bruckenau and Bocklet (cold).
6. Tunbridge (cold).
7. Niton (cold).
8. Furnas (hot).

VIII. Ioduretted and brominated.

1. Creuznach.
2. Hall.
3. Ischl.
4. Woodhall.

WHITE SULPHUR.

	In a wine gallon.
Sulphuretted Hydrogen,.....	cubic inches. 2·5
Carbonic Acid,.....	2
Oxygen,.....	1·448
Nitrogen,.....	3·552
Total,.....	9·5

	Solid contents in a pint.
Sulphate of Magnesia,.....	grs. 5·588
Sulphate of Lime,.....	7·744
Carbonate of Lime,.....	1·150
Chloride of Calcium,.....	0·204
Chloride of Sodium,.....	0·180
Oxide of Iron,.....	a trace.
Loss,.....	0·410
Total,.....	15·276

PROF. W. B. ROGERS.

BALSTON.

SANS SOUCI SPRING.

	In a wine gallon.
Chloride of Sodium,.....	grs. 143·733
Bicarbonate of Soda,.....	12·66
Bicarbonate of Magnesia,.....	39·1
Carbonate of Lime,.....	43·407
Carbonate of Iron,.....	5·95
Iodide of Sodium,.....	1·3
Silica,.....	1
Total,.....	247·15

STEEL.

SARATOGA.

IODINE SPRING.

	Gaseous contents in a wine gallon.
Carbonic Acid,.....	cubic inches. 336
Atmospheric Air,.....	4
	Total, 340

	Solid contents in a wine gallon.
Chloride of Sodium,.....	grs. 187
Carbonate of Magnesia,.....	75
Carbonate of Lime,.....	26
Carbonate of Soda,.....	2
Carbonate of Iron,.....	1
Iodine,.....	3·5
	Total, 294·5

PROF. EMMONS.

SARATOGA.

PAVILION SPRING.

	Gaseous contents in a wine gallon.
Carbonic Acid,.....	cubic inches. 359·05
Atmospheric Air,.....	5·03
	Total, 364·08

	Solid contents in a wine gallon.
Chloride of Sodium,.....	grs. 187·68
Carbonate of Soda,.....	4·92
Carbonate of Lime,.....	52·84
Carbonate of Magnesia,.....	56·92
Carbonate of Iron,.....	3·51
Sulphate of Soda,.....	1·48
Iodide of Sodium,.....	2·59
Alumina,.....	0·42
Silica,.....	1·16
Phosphate of Lime,.....	0·19
Bromide of Potassium,.....	a trace.

Total, 311·71

DR. J. R. CHILTON.

UNION SPRING.

	Gaseous contents in a wine gallon.
Carbonic Acid,.....	cubic inches. 314·16
Atmospheric Air,.....	4·62
	<hr/>
	Total, 318·78

	Solid contents in a wine gallon.
Chloride of Sodium.....	grs. 243 620
Carbonate of Magnesia,.....	84 265
Carbonate of Lime,.....	41·600
Carbonate of Soda,.....	12 800
Carbonate of Iron,.....	5·452
Iodide of Sodium, or Iodine,.....	3 600
Silica and Alumina,.....	1·570
Bromide of Potassium,.....	a trace.
	<hr/>
	Total, 392·907

DR. J. R. CHILTON.

CONGRESS SPRING.

	Gaseous contents in a wine gallon.
Carbonic Acid,.....	cubic inches. 311
Atmospheric Air,.....	7
	<hr/>
	Total, 318

	Solid contents in a wine gallon.
Chloride of Sodium,.....	grs. 385
Iodide of Sodium,.....	3·5
Bicarbonate of Soda,.....	8·982
Bicarbonate of Magnesia,.....	95·788
Carbonate of Lime,.....	98·098
Carbonate of Iron,.....	5 075
Silica,.....	1·5
Bromide of Potassium,.....	a trace.
	<hr/>
	Total, 597·943

STEEL.

VI. SALINE DRAUGHTS.

Proportional quantities of Alkalies and Acids required for saturation.

ALKALINE CARBONATES.	ACIDS.		
	Lemon Juice.	Citric Acid.	Tartaric Acid.
20 Grains.	Drachms.	Grains.	Grains.
Potassæ bicarbonas,*.....	3½	14	15
— carbonas (<i>olim</i> subcarbonas),*.....}	4	17	18
Sodæ sesquicarbonas,.....	4	17	18
— carb. (<i>olim</i> subc.)†.....	2½	9½	10½
Ammoniæ sesquic. (The only salt in use.).....}	6	24	25½

* When tartaric acid is employed it is necessary to add the acid to the potash solution with care; otherwise, when added in excess, the insoluble bitartrate forms.

† Crystallized; 16 grs. of this, when dried, yield 6 grs. of sod. carb. exsicc.

VII. Weights and Measures. (See pp. 15, 16.)

INDEX.

PAGE		PAGE	
Absinthium	36	Alumen	39
Acacia	52	exsiccatum	122
Acetosella	90	Ammonia	18
Acetum	89	Ammoniacum	28
colchici	79	Ammoniæ sesquicarbonas	18-49-61
cantharidis	111	Amygdala amara	52
destillatum	89	dulcis	ib.
scillæ	59	Amylum	53
Acidum benzoicum	29	Anethum	45
aceticum	89-111	Anisum	ib.
arsenicum	101	Anthemis	36-61
eiticum	89	Antimonii pot. tart.	61-65
hydrochloricum	101	Aqua anethi	45
hyd. dilutum	89-101	aurantii	37
nitrie. dilutum	89	carui	46
hydroey. dilut.	67	cinnamomi	33
phospho. dilut.	102	fœniculi	46
sulph. dilut.	89-102	menthæ piperitæ	47
tartaricum	89	pulegii	48
Aconitina	118	viridis	ib.
Aconitum	68	pimentæ	35
napellus	127	Argenti nitras	102-111
Acorus	36	Armoracia	94
Acrid poisons	128	Asarum	62
Æther sulphuricus	27	Aspidium	26
Alcohol	93	Assafetida	28
Allium	65-110	Aurantii cortex	36
Aloes	76	Aurantium	37
extractum	ib.	Avena	53
Althæa	52		

	PAGE		PAGE
Balsamum peruvianum.	65	Ceratum hydrargyri com-	
tolutanum ..	ib.	posit.	114
Barii chloridum	21	plumbi acetatis	113
Baths	128	comp.	ib.
Belladonna	69	resinæ	ib.
emplastrum.....	ib.	sabinæ	112
extractum	ib.	saponis	113
Benzoinum	28	Cetaceum	53
Bismuthi trisnitras	103	Cetraria	ib.
Borax	40	Chimaphila	56
Brominium	21	Cinchona	106
Cajuputi oleum	95	Cinnamomum	33
Calamina præparata	122	Colchicum	78
Calumba	105	Colocynthis	79
Calx	18	Confectio amygdalæ	52
Cambogia	78	aromatica	33
Camphora	69	aurantii	37
Canella	31	cassiaæ	78
Cantharis	55	opii	73
Capsicum	31-112	piperis nigri	35
Cardamine	29	rosæ caninæ	91
Cardamomum	32	gallicæ	43
Carui	46	rutæ	64
Caryophyllus	32	scammonii	86
Caryophylli oleum	ib.	sennæ	ib.
Cascarilla	37	Conium	70
Cassia	78	Contrajervia	33
Castoreum	29	Copaiba	66
Cataplasma allii	110	Coriandrum	46
aluminis	ib.	Corrosive and irritant	
carbonis ligni	111	poisons	123
conii	110	Counterpoisons	ib.
dauci	111	Creosoton	95-113
fermenti	110	Creta præparata	18-122
lini	55-110	Crocus	29
sinapis	35-110	Cubeba	34
Catechu	40	tinctura	ib.
Centaurium	106	Cupri ammonio-sulphas	29
Ceratum	113	sulphas	41-62-103
calaminæ	ib.	Cuprum	41
cantharidis	111	Cusparia	38
cetacei	113	Cydonia	53

	PAGE		PAGE
Cyminum	46	Enema emoliens	117
Daucus	ib.	fætidum	ib.
Decoctum aloës comp.	77	opii	ib.
amyli	53	rutæ	ib.
cetrariae	ib.	tabaci	ib.
chimaphilæ	56	terebinthinæ	88-117
cinchonæ	106-118	Ergota	63
cydoniæ	53	Euphorbium	112
dulcamaræ	50-57	Extractum aconiti	68
granati	41	Extr. aloës purificatum	76
hordei	54	cinchonæ	107
comp.	ib.	colchici aceticum ..	79
malvæ com.	55-120	cormi ..	ib.
papaveris	120	colocynthidis	80
quercus	43-118-121	compositum ..	ib.
sarzæ	25	conii....	70
compositum ...	ib.	digitalis	56
scoparii compositum	59	elaterii....	80
senegæ	ib.	gentianæ	108
tormentillæ	44	glycyrrhizæ....	54
ulmi	ib.	hæmatoxyli	42
uvæ ursi	44-61	hyoscyami	71
veratri	121	jalapæ	81
Digitalis	56	lactucæ	71
Diosma	50	lupuli	39
Doses , proportion of, according to age	130	opii purificatum	72
Dulcamara	50-57	papaveris	74
Elaterium	80	pareiræ	58
Emplastrum ammoniaci	116	rhei	85
cum hydrarg.	ib.	sarzæ	25
belladonnae	ib.	stramonii....	75
cantharidis	111	taraxaci	60
cere	116	uvæ ursi	45-61
galbani	ib.	 Ferri ammon.-chloridum	103
hydrargyri	ib.	iodidum	25-104
opii	ib.	potassio tartras....	105
plumbi	ib.	sesquioxidum	103
saponis	ib.	sulphas	104
Enëma aloës	78-117	Ficus	54
catharticum	117	Fœniculum	46
colocynthidis....	80-117	Gallæ	41
		Galbanum	29

	PAGE		PAGE
Gentiana	108	Infusum lini comp.	54
Glycyrrhiza	54	Inula	34
Granatum	26-41	Iodinum	24
Guaiacum.....	50	Ipecacuanha	50-62
Hæmatoxylum.....	42	Kino	42
Helleborus	80	Krameria	ib.
Hordeum.....	54	Lactucæ extractum	71
Hydrargyri bichloridum	22	Lavandula	46
biniodidum.....	23	Laurus	47
chloridum.....	22-81-122	Limones	38
iodidum.....	22	Linimentum ammoniæ	119
nitrico-oxidum.....	122	ammon. sesquicarb.	ib.
oxidum	22	compositum	ib.
sulph. cum sulphure	23	calcis	121
ung. cerat. &c. ..	114-115	camphoræ	119
compositum	ib.	compositum	ib.
Hydrargyrum	21-80	crotonis	120
cum creta.....	80	æruginis	121
Hyoscyamus.....	71	hydrargyri comp.	119
Jalapa	81	opii.....	120
Juniperus	57	saponis	119
Infusum anthemidis... 36-61		terebinthinæ	120
armoracie composit. 94		Linum	54
aurantii composit.. 37		Liquor alumin. comp. 40-121	
calumbæ	105	ammoniæ	119
caryophylli	32	acetatis	49-90
cascarillæ	38	sesquicarbonatis 18	
catechu comp. ... 40-118		barii chloridi	21-102
cinchonæ	106-118	calcii chlorid	21-103
cuspariæ	38	calcis	18-121
digitalis.....	56	cupri ammon.-sulph. 121	
diosmæ	50	hydrarg. bichloridi ..	22
gentianæ composit..	108	liniment. ammoniæ	119
lupuli.....	38	plumbi diacetatis ..	120
quassiaæ	109	dilutus	ib.
krameriaæ	42	potassæ	19
pareiræ	57	effervescens	20
rhei.....	84	arsenitis	101
rosæ comp. 43-118		carbonatis	20-58
scoparii	59	potassii iodidi comp.	24
sennæ compositum ..	86	sodæ effervescens	20
serpentariaæ	95	Lobelia	62
simarubæ	109		

	PAGE		PAGE
<i>Lotio ammoniæ hydroc.</i>	120	<i>Oleum amygdalæ</i>	53
<i>flava</i>	121	<i>aurantii</i>	37
<i>hydrocyanica</i>	ib.	<i>anisi</i>	45
<i>nigra</i>	ib.	<i>anthemidis</i>	36
<i>Lupulus</i>	38	<i>cajaputi</i>	120
<i>Magnesia</i>	19-82	<i>carui</i> ,	46
<i>Magnesiæ carbonas</i>	19	<i>caryophylli</i>	32
<i>sulphas</i>	82	<i>cinnamomi</i>	33
<i>Malva</i>	55	<i>juniperi</i>	57
<i>Manna</i>	82	<i>lavandulæ</i>	46
<i>Marrubium</i>	66	<i>menthæ piperitæ</i>	47
<i>Mastiche</i>	66	<i>pulegii</i>	48
<i>Mel boracis</i>	40-118	<i>viridis</i>	ib.
<i>rosæ</i>	44	<i>myristicæ</i>	34
<i>Mentha piperita</i>	47	<i>origani</i>	48
<i>pulegium</i>	ib.	<i>pimentæ</i>	34
<i>viridis</i>	48	<i>ricini</i>	85
<i>Menyanthes</i>	108	<i>rosmarini</i>	48
<i>Mezereon</i>	112	<i>succini</i>	31
<i>Mezereum</i>	51	<i>terebinthinæ</i>	26-61-88
<i>Mineral waters, classified</i>		<i>tiglii</i>	88
list of	131	<i>Olibanum</i>	30
<i>Mistura acaciæ</i>	52	<i>Opium</i>	72
<i>ammoniaci</i>	28	<i>Opopanax</i>	30
<i>amygdalæ</i>	52	<i>Origanum</i>	48
<i>assaftidæ</i>	28	<i>Oxymel</i>	55
<i>camphoræ</i>	69	<i>scillæ</i>	59
<i>cascarillæ composita</i>	38	<i>Papaver</i>	74
<i>cretæ</i>	19	<i>Pareira</i>	57
<i>ferri composita</i>	104	<i>Petroleum</i>	30
<i>gentianæ composita</i>	108	<i>Pilulæ aloës compositæ</i>	76
<i>guaiaci</i>	50	<i>cum myrrhâ</i>	ib.
<i>moschi</i>	30	<i>cambogiæ comp.</i>	77-78
<i>spiritus vini gallici</i>	97	<i>conii compositæ</i>	70
<i>Mora</i>	90	<i>ferri compositæ</i>	105
<i>Morphiæ acetas</i>	73	<i>galbani compositæ</i>	29-31
<i>hydrochloras</i>	ib.	<i>hydrargyri</i>	22-31
<i>Moschus</i>	30	<i>iodidi</i>	23
<i>Mucuna</i>	26	<i>chloridi comp.</i>	22
<i>Myristica</i>	34	<i>ipecac. compositæ</i>	51
<i>Myrrha</i>	66	<i>rhei compositæ</i>	85
<i>Nux vomica</i>	71	<i>saponis compositæ</i>	72

	PAGE		PAGE
Pilulæ sagapeni comp..	31-77	Rumex	91
scillæ compositæ	59	Ruta	64-112
Pimenta.....	34	Sabina	64
Piper longum	35	Sagapenum	31
nigrum.....	35-112	Saline draughts.....	135
Plumbi acetns	43	Sapo	20
iodidum.....	24	Sarza	25
Plumbum	43	Sassafras	51
Poisons	123	Scammonium.....	85
Potassæ acetas	58-82-91	Scilla	58
bicarbonas.....	20-58	Scoparius	59
bisulphas.....	83	Senega	51
bitartras.....	83-91	Senna	86
carbonas	20-58	Serpentaria	95
nitrás	58	Simaruba	109
sulphas	83	Sinapis	35-63
tartras	ib.	Sodæ acetas	87
hydras	111	carbonas	20
Potassa cum calce.....	ib.	exsiccata.....	ib.
Potassii bromidum	21	phosphas	87
iodidum	24	potassio-tartras	ib.
Pruna	84	sesquicarbonas	20
Pulvis aloës compositus	76	sulphas	87
antimonii comp.	49	Spigelia	27
cinnamomi comp.	33	Spiritus ætheris nitrici..	60
cretæ compositus....	18	sulph. comp....	93
cum opio	ib.	ammoniæ aromaticus	94
jalapæ compositus..	81	fætidus.....	ib.
ipecacuanhæ comp.	51	anisi	45
kino compositus....	42	armoraciæ composit.	94
scammonii comp.	86	carui	46
tragacanthæ comp.	55	cinnamomi	33
Quassia	109	juniperi compositus	57
Quercus	43	lavandulæ	47
Quinæ disulphas	107	menthæ piperitæ ...	ib.
Rhamnus	84	menthæ pulegii....	48
Rheum	ib.	viridis	ib.
Rhei pulvís	122	myristicæ	34
Rosa canina	91	pimentæ	ib.
centifolia	85	rectificatus	93
gallica	43	rosmarini	49
Rosmarinus	48	tenuior.....	93

	PAGE		PAGE
Spiritus vini gallici	96	Tinctura cinchonæ	106
Stannum.....	27	composita	107
Stramonium	75	cinnamomi	33
Strychnia	72	composita	ib.
Succinum	31	colchici	79
Sulphur	51	composita	ib.
Syrupus althææ.....	52	conii	70
aurantii.....	37	cubebæ	34
mori	90	digitalis.....	56
papaveris	74	ferri ammon.-chlor..	104
rhamni	84	sesquichloridi..	ib.
rosæ	85	gallæ.....	41-118
sarzæ	25	gentianæ compositæ	108
sennæ	87	guaiaci	50
tolutanus	66	composita	ib.
zinziberis	35	hellebori	80
Tabacum	63	hyoscyami	71
Tamarindus	88	iodini comp.	24
Taraxacum	60	jalapæ	82
Tela vesicatoria	111	kino.....	42-118
Terebinthina	60	lavandulæ composita	47
canadensis	ib.	lupuli.....	39
chia	61	myrrhæ	66-118
vulgaris	60	opii	72
Testæ præparatae	19	rhei composita.....	84
Tiglii oleum	88	scillæ	59
Tinctura aloës	77	sennæ composita...	86
Tinctura aloës comp....	ib.	serpentariae	95
ammoniæ composita	94	valerianæ	31
assafetidiæ	28	composita	ib.
aurantii	37	zinziberis	35
balsami tolutani....	66	Tomentilla	44
benzoini composita.	29	Toxicodendron	75
calumbæ	106	Tragacantha	55
camphoræ composita	70	Unguentum ant. pot. tar.	112
cantharidis	56	aconitinæ	115
capsici.....	32-118	belladonnæ	ib.
cardamomi	32	cantharidis	111
composita	ib.	cetacei	113
cascarillæ	38	creosoti	ib.
castorei	29	elemi	114
catechu	40	gallæ comp.....	41-114

	PAGE		PAGE
Unguentum hydrargyri		Unguentum veratri.....	ib.
ammonio.....	114	veratriæ	ib.
biniodidi	115	zinci.....	113
fortius	114	Uva ursi	61
iodidi	115	Valeriana	31
iodini	ib.	Veratrica	75
compositum	ib.	Veratrump.....	63-75
mitius	114	Vinum aloes	78
nitratis.....	ib.	antimonii pot. tart..	61
nitrico-oxidi...	ib.	colchici	79
mezerei	112	ipecacunhæ.....	51-62
picis liquidи	114	opii	72
nigri.....	ib.	veratri	63
plumbi compositum	ib.	Weights and measures..	15
iodidi.....	115	Zinci oxydum	105
sambuci	113	acetas	45
sulphuris	115	sulphas.....	45-63-105
comp.	ib.	Zinziber	35

M E D I C A L B O O K S ,

PUBLISHED BY

SAMUEL S. & WILLIAM WOOD,

261 Pearl-street,

NEW YORK.

THE FIRST LINES OF THE THEORY AND PRACTICE OF SURGERY ; including the principal operations. By SAMUEL COOPER, Professor of Surgery in University College Hospital, London, &c. With notes and additions by WILLARD PARKER, M.D., Professor of Surgery in the College of Physicians and Surgeons, in the University of the State of New York, &c. 4th American, from the 7th London, edition.

"The work of Samuel Cooper, having passed through *seven* English, and *four* American, editions, requires no commendation to ensure the present edition an early sale. It abounds in accurate principles and practical precepts, the result of extensive experience and careful observation ;" and the additions by Prof. Parker, "are highly judicious, and in no small degree enhance the value of the original work."—*Western Lancet.*

THE INFLUENCE OF TROPICAL CLIMATES ON EUROPEAN CONSTITUTIONS. By JAMES JOHNSON, M.D., and JAMES RANALD MARTIN, Esq. From the sixth London edition, with Notes by an American Physician.

"This, *the Book* on Tropical Climates, scarcely needs commendation at our hands, and there is little in its many pages except the promulgation of sound truths, and as sound deductions. Almost every American medical man has read and been benefited by this early production of the lamented author; an author who has done more through his numerous works, with the '*Medical and Chirurgical Journal*,' to mould correctly the medical opinion in these United States, than even any, or many of its eminent native writers.

"We would advise all who do not possess this work, and who value information and amusement, to obtain it."—*New York Journal of Medicine*.

"Of the comparatively few medical books, amongst the many with which the press teems, destined to survive the changes incident to progressive advancement in science, the celebrated work on Tropical Climates, by Dr. Johnson, the late distinguished editor of the *Medico-Chirurgical Review*, is a prominent example."—*Buffalo Medical Journal*.

THE ANATOMICAL REMEMBRANCER; or, Complete Pocket Anatomist: containing a concise description of the Bones, Ligaments, Muscles, and Viscera, the distribution of the Nerves, Blood-Vessels, and Absorbents, the ar- rangement of the several Fasciæ, the Organs of Generation in the Male and Female, and the Organs of the Senses.

"It contains but two hundred and fifty pages, and is really an anatomical *multum in parvo*."—*London Med. & Surg. Journal*.

"Every medical man, as well as pupil, would do well to have it on his table."—*Buffalo Medical Journal*.

"It will be found of great assistance to those who frequent the dissecting room, or are desirous of renewing at a glance, the knowledge they have acquired at some former period."—*Southern Journal of Medicine and Pharmacy*.

A MANUAL OF CHEMISTRY. By RICHARD D. HOBLYN, A.M., OXON.

"It is an excellent work of its kind, and the author enters as fully into the subject as he can, and indeed as fully as is

necessary in a manual 'prepared for the purpose of affording public or private instructions in the principles of chemistry.' He has stated 'the most important facts and theories of the science in as condensed a form as the subject admits.'—"New York Journal of Medicine.

"In descriptions somewhat parenthetical, yet clear, the author contrives to communicate a great deal of information to the learner in a small space."—*Bulletin of Med. Science.*

"It is rarely that we meet with a manual so well adapted for the object of its publication.

"The most important facts and theories of the science are stated in as condensed a form as the subject admits.

"The work is tersely written, and with so much conciseness, that a great amount of matter is comprised in comparatively a few pages. We know, indeed, of no work on the same subject, that contains so clear and comprehensive an account of the principles of chemistry, within any thing like so brief a space."—*Medical Examiner.*

LECTURES ON NATURAL AND DIFFICULT PARTURITION. By EDWARD WILLIAM MURPHY, A.M., M.D., Professor of Midwifery, University College, London.

"They are characterized by a profound knowledge of the subjects on which they treat, and attest the careful, thoughtful, modest, and discriminating mind of the author."—*Med. Chir. Review.*

"The student will find it replete with accurate and lucid instruction."—*London Medical Gazette.*

THE MEDICAL REMEMBRANCER ; or, Book of Emergencies ; in which are concisely pointed out the immediate remedies to be adopted in the first moments of danger from Poisoning, Drowning, Apoplexy, Burns and other accidents, with the tests for the principal Poisons, and other useful information. By EDWARD B. L. SHAW, M.R.C.S., and L.A.S., &c. Revised and Improved by an American Physician.

THE YOUNG STETHOSCOPIST ; or, The Student's Aid to Auscultation. By HENRY J. BOWDITCH, M.D.

"Plain and simple in its arrangement, accurate in details, and sufficiently extensive for elementary purposes, it should become the companion of every student of auscultation."—*Western Lancet.*

A TREATISE ON DISEASES OF THE CHEST, AND ON MEDIATE AUSCULTATION. By R. T. H. LAENNEC, M.D. Translated from the 3d French edition, with copious notes, a Sketch of the Author's Life, and an extensive bibliography of the different diseases. By JOHN FORBES, M.D., F.R.S. To which are added the Notes of Professor Andral, contained in the 4th and latest French edition, translated and accompanied with observations on Cerebral Auscultation. By JOHN D. FISHER, M.D. With Plates.

"We cannot occupy more of our space at present with further extracts from this most valuable work; but we cannot conclude without recommending it to our readers in the strongest terms, and *for the third time*, but, we hope not the last, as the most comprehensive and valuable work which has ever been published on the pathology and diagnosis of diseases of the chest."—*Med. Chir. Review.*

AN ESSAY ON INDIGESTION ; or, Morbid Sensibility of the Stomach and Bowels, as the Proximate Cause or Characteristic Condition of Dyspepsy, Nervous Irritability, Mental Despondency, Hypochondriasis, and many other Ailments of Body and Mind. To which are added observations on the Diseases and regimen of Invalids on their return from hot and unhealthy climates. By JAMES JOHNSON, M.D., Editor of Med. Chir. Review, &c.

S. S. & W. Wood have recently published

A BEAUTIFUL

LITHOGRAPHIC PLATE,

Thirty by twenty-four inches, containing the *Portraits* of the *Medical Faculty of the University of New York*, viz.—Chancellor FRELINGHUYSEN, and Professors MOTT, PATTISON, REVERE, PAINE, BEDFORD, and DRAPER.

Also, one of the same size,

Containing the *Portraits* of the *Faculty of the College of Physicians and Surgeons of the University of the State of New-York*, viz.—Professors STEVENS, SMITH, BECK, TORREY, WATTS, PARKER, and GILMAN.

